


# LAKE WENTWORTH

## Water Quality Monitoring: 2002

### Summary and Recommendations

#### NH LAKES LAY MONITORING PROGRAM



By: Robert Craycraft & Jeffrey Schloss

Center for Freshwater Biology  
University of New Hampshire




UNIVERSITY of NEW HAMPSHIRE  
COOPERATIVE EXTENSION

To obtain additional information on the NH Lakes Lay Monitoring Program (NH LLMP) contact the Coordinator (Jeff Schloss) at 603-862-3848 or Assistant Coordinator (Bob Craycraft) at 603-862-3696.

# PARAMETERS SAMPLED

## NH LAKES LAY MONITORING PROGRAM



Center for Freshwater Biology (CFB) corroboration with the lay monitor data includes assessment of 1) physical parameters (water transparency, temperature profiles, light transmission profiles and water color); 2) chemical parameters (dissolved oxygen profiles, "free" carbon dioxide, total alkalinity, pH, total phosphorus and specific conductivity profiles); 3) biological parameters (chlorophyll a, phytoplankton community and zooplankton community). Note: in addition to the above parameters, other measurements are often collected at the discretion of the CFB or at the request of the lake association.

# PREFACE

---

This report contains the findings of a water quality survey of Lake Wentworth and Crescent Lake, Wolfeboro New Hampshire, conducted in the summer of 2002 by the University of New Hampshire **Center for Freshwater Biology (CFB)** in conjunction with the Lake Wentworth Association.

The report is written with the concerned lake resident in mind and contains a brief, non-technical summary of the year 2002 results as well as more detailed "Introduction" and "Discussion" sections. Graphic display of data is included, in addition to listings of data in appendices, to aid visual perspective.

# ACKNOWLEDGMENTS

The year 2002 was the nineteenth year the Lake Wentworth Association participated in the **New Hampshire Lakes Lay Monitoring Program (LLMP)**. The volunteer monitors involved in the water quality monitoring effort are highlighted in Table 1 while John Nichols again coordinated the volunteer monitoring activities on Lake Wentworth and Crescent Lake and acted as liaison to the **Center for Freshwater Biology (CFB)**. The **CFB** congratulates the volunteer monitors on the quality of their work, and the time and effort put forth. We invite other interested residents to join the Lake Wentworth water quality monitoring effort in 2003 and expand upon the growing database. Funding for the water quality monitoring program was provided by the Lake Wentworth Association.

The **Center for Freshwater Biology** is a not-for-profit research program coordinated by Jeffrey Schloss and Robert Craycraft. Members of the **CFB** summer field team included, Autumn Carlson, William Clark, Robert Craycraft, Juliette Nowak and Kirsten Pulkkinen while Lydia Pitkin, Melissa McCartney, Santhana Souksamrane and Amy Surprenant provided additional laboratory support in the fall.

The **CFB** acknowledges the University of New Hampshire Cooperative Extension for funding and furnishing office and storage space while the College of Life Sciences and Agriculture provided laboratory facilities and additional storage space. A generous gift from the **Samuel P. Pardoe Foundation** allowed for an update of our volunteer's temperature profiling equipment and the **US Geological Survey**, through the **UNH Water Resources Research Center**, provided some resources for staff support.

Participating groups in the **LLMP** include: The Center Harbor Bay Conservation Commission, Dublin Garden Club, Eaton Conservation Commission, Governor's Island Club Inc., Laconia Conservation Commission, Meredith Bay Rotary Club, The New Hampshire Audubon Society, North River Lake Water Quality Audit Committee, Society for Protection of Lakes and Streams, Walker's Pond Conservation Society, United Associations of Alton, the associations of Baboosic Lake, Berry Bay, Bow Lake Camp Owners, Chalk Pond, Lake Chocorua, Cunningham Pond, Crystal Lake, Dublin Lake, Goose Pond, Great East Lake, Langdon Cove, Long Is-

**Table 1: Lake Wentworth and Crescent Lake Volunteer Monitors (2002)**

Monitor Name
John Nichols
Brennan Conley
Alan Thompson
Liz Thompson
Garrett Conley
Steven Rohde
Terry Conley
Dana Bartlett
David McLarnon
Ken Roberts
Laura Nichols
Lioudmila Mojaiskaia
Ryan McCarthy
B. Sandford
Gary Baker
Nate Roberts
Barbara Baker
Joe Anderer
David Barnes
Don Kretchmer
Jim Hulm

land Landowners, Lovell Lake, March's Pond, Mendum's Pond, Merrymeeting Lake, Milton Ponds Lake Lay Monitoring, Mirror Lake (Tuftonboro), Moultonborough Bay, Lake Winnepesaukee, Naticook Lake, Newfound Lake, Nippo Lake, Silver Lake (Madison), Silver Lake (Tilton), Squam Lakes, Sunset Lake, Wentworth Lake and the towns of Alton, Amherst, Enfield, Errol, Madison, Meredith, Merrimack, Milan, Strafford and Wolfeboro.

# Crescent Lake

## 2002 Non-Technical Summary

---

Water quality data were collected by the Crescent Lake volunteer monitors between June 16 and September 20, 2002. A more in depth water quality survey of the Crescent Lake deep sampling station (Site 6 Center) was conducted by the **Center for Freshwater Biology (CFB)** on August 1, 2002 to augment the volunteer monitoring data. Generally speaking, the year 2002 Crescent Lake water quality remained high as characterized by the high water clarity, the low levels of microscopic plant "algal" growth and low phosphorus, nutrient, levels (Table 2).

For the second consecutive year, many participating New Hampshire Lakes Lay Monitoring Program lakes were characterized by near record/record high water transparency measurements. Furthermore, the 2002 microscopic plant "algal" levels included some of the lower concentrations documented over the past ten years of water quality monitoring for many participating lakes that included Crescent Lake (Figure 22). The dry summer months limited the flushing of sediments and nutrients into our lakes; the same nutrients that make our lawns green will also stimulate plant growth in our New Hampshire lakes. Thus, the low levels of microscopic plant growth and the very clear waters that characterized most New Hampshire lakes are associated (at least in part) with the limited summer 2002 rainfall.

The following section discusses the year 2002 and historical Crescent Lake water quality data. *Refer to Appendix A for a complete listing of the year 2002 Crescent Lake water quality data.*

**Table 2: 2002 Crescent Lake Seasonal Average Water Quality Readings and Water Quality Classification Criteria used by the New Hampshire Lakes Lay Monitoring Program.**

Parameter	Oligotrophic "Pristine"	Mesotrophic "Transitional"	Eutrophic "Enriched"	Crescent Lake Average (range)	Crescent Lake Classification
<b>Water Clarity (meters)</b>	> 4.0	2.5 - 4.0	< 2.5	5.3 meters (range: 4.7 - 5.9)	Oligotrophic
<b>Chlorophyll a (ppb)</b>	< 3.0	3.0 - 7.0	> 7.0	1.6 ppb (range: 0.7 - 2.4)	Oligotrophic
<b>Phosphorus (ppb)</b>	< 15.0	15.0 - 25.0	> 25.0	* 8.1 ppb (range: 7.7 - 8.4)	Oligotrophic

\* Total Phosphorus data were collected by the Center for Freshwater Biology on August 1, 2002).

1) **Water Clarity (measured as Secchi Disk transparency)** – The year 2002 Crescent Lake water clarity values were high and remained well above the transparency of 13.2 feet (4 meters) that is considered the boundary between an unproductive "pristine" and more nutrient enriched "transitional" New Hampshire lake (Figures 13 & 14).

A Comparison between the year 2002 and the historical (1981-2001) water transparency data indicate the seasonal average Secchi Disk transparency values increased (i.e. clearer water) for the third consecutive year (Figure 21). *Note: the Crescent Lake sampling site is shallow and the Secchi Disk sometimes*

reaches the lakebottom before disappearing from view. Thus, the Secchi Disk measurements should be interpreted with caution.

2) **Microscopic plant abundance “greenness” (measured as chlorophyll  $\alpha$ )** – All of the 2002 Crescent Lake chlorophyll  $\alpha$  concentrations remained below the concentration of 3 parts per billion (ppb) that is considered the boundary between a nutrient poor and more nutrient enriched “greener” New Hampshire lake (Figure 13).

The year 2002 seasonal chlorophyll  $\alpha$  concentration documented in Crescent Lake is one of the lower values (i.e. less algal greenness) documented since volunteer monitoring was instituted on Crescent Lake during the summer of 1984 (Figure 22). No new maximum or minimum chlorophyll  $\alpha$  concentrations were documented in Crescent Lake during the 2002 sampling season.

3) **Background (dissolved) water color : often perceived as a “tea” color in more highly stained lakes** – The year 2002 Crescent Lake dissolved color concentration averaged 11.9 chloroplatinate units (cpu) and fell within the classification of a slightly “tea” colored lake (Table 3). Dissolved color, or true color as it is sometimes called, is indicative of dissolved organic carbon levels in the water (a by-product of microbial decomposition). Small increases in water color from the natural breakdown of plant materials in and around a lake are not considered to be detrimental to water quality. However, increased color can lower water transparency, and hence, change the public perception of water quality.

**Table 3. Dissolved Color Classification Criteria used by the New Hampshire Lakes Lay Monitoring Program.**

Range	Classification
0 - 10	Clear
10 - 20	Slightly colored
20 - 40	light tea color
40 - 80	tea colored
> 80	highly tea colored

4) **Total Phosphorus: the nutrient considered most responsible for elevated microscopic plant growth in our New Hampshire Lakes.** - Total phosphorus concentrations were low in Crescent Lake when measured on August 1, 2002. The Crescent Lake total phosphorus concentrations ranged from 7.7 to 8.4 parts per billion (ppb) and remained below the concentration of 10 ppb that is considered sufficient to cause an algal bloom.

5) **Resistance against acid precipitation (measured as total alkalinity)** – The year 2002 Crescent Lake alkalinity of 7.1 milligrams per liter (mg/l) is characteristic of a lake with a moderate vulnerability to acid precipitation according to the standards devised by the New Hampshire Department of Environmental Services (Table 4). Generally speaking, the geology of the

**Table 4. Alkalinity Classification Criteria used by the New Hampshire Department of Environmental Services**

Range	Classification
< 0	Acidified
0 - 2	Extremely Vulnerable
2.1 - 10.0	Moderately Vulnerable
10.1 - 25.0	Low Vulnerability
> 25.0	Not Vulnerable

region does not contain the mineral content (e.g. limestone) which increases the buffering capacity in our surface waters. Thus, lakes in the vicinity (i.e. Lake Winnepesaukee and Merrymeeting Lake) have naturally low alkalinities.

**Lake acidity (measured as pH)** - The year 2002 Crescent Lake pH data, collected by the **Center for Freshwater Biology** on August 1, 2002, ranged from 6.7 to 6.8 units and remained well within the tolerable range for most aquatic organisms.

6) **Dissolved salts: measured as specific conductivity** - Specific Conductivity levels, documented in Crescent Lake, were low to moderate and ranged from 59.0 to 63.0 micro-Siemans ( $\mu$ S) when measured on August 1, 2002. High specific conductivity values can be an indication of problem areas around a lake where failing septic systems, heavy fertilizer applications and sedimentation contribute "excessive" nutrients that make their way into Crescent Lake. High specific conductivity values can also be associated with road salt runoff that is flushed into our New Hampshire lakes.

7) **Temperature and dissolved oxygen profiles** - Temperature profiles collected by the volunteer monitors indicate Crescent Lake only becomes partially stratified during the summer months. During relatively calm periods, Crescent Lake develops a warm upper water layer, the **epilimnion**, which overlies a layer of rapidly decreasing temperatures, the **thermocline**. The formation of thermal stratification limits the replenishment of oxygen in the deeper waters and under adverse conditions can result in oxygen depletion near the lake-bottom.

**Dissolved oxygen concentrations required for a healthy fishery** - Dissolved oxygen concentrations documented by the **Center for Freshwater Biology** on August 1, 2002 remained high and above the concentration of 3 milligrams per liter down to the lakebottom (Figure 29). The dissolved oxygen concentration of 3 milligrams per liter is commonly considered the minimum dissolved oxygen concentration required for the successful growth and reproduction of most warmwater fish species that include bass and perch.

8) A Comparison between the **Center for Freshwater Biology Group** and lay monitor data indicate the Crescent Lake volunteer monitors are doing an excellent job of collecting water quality data.

9) Based on the current and historical water quality data, Crescent Lake would be considered an unproductive "pristine" lake that at times exhibits some characteristics (i.e. reduced water transparency and short-term algal blooms) that are typical of a more nutrient enriched "transitional" lake. A first step towards preserving high water quality characteristic of Crescent Lake is to take action at the local level and do your part to minimize the number of pollutants (particularly sediment and the nutrient phosphorus) that enter the lake. Whenever possible, **maintain riparian buffers** (vegetative buffers adjacent to the water body). These buffers will biologically "take up" nutrients before the nutrients enter the lake and will also provide physical filters that allow materials to



settle out before reaching the lake. **Reduce fertilizer applications.** Most residents apply far more fertilizers than necessary which can be a costly expense to the homeowner and can also be detrimental to the lake since the same nutrients that make our lawns green will also stimulate plant growth in our lakes. **Make sure your septic system is well maintained.** Have your septic system pumped out on a regular basis. An improperly functioning septic system can contribute "excessive" nutrients into the lake and result in early failure, costing thousands of dollars to repair or replace. Future volunteer monitoring efforts should be directed at pinpointing problematic regions around the lake where corrective and educational efforts should be focused. It is important to make sure the watershed residents are well-educated on water quality related issues. Numerous publications are available through University of New Hampshire Cooperative Extension, the New Hampshire Lakes Association, the New Hampshire Department of Environmental Services as well as several other local, state and federal agencies. It is imperative that future activities within the Crescent Lake watershed are carefully thought out before implementation if water quality degradation is to be minimized.

# Lake Wentworth

## 2002 Non-Technical Summary

---

Water quality data were collected by the Lake Wentworth volunteer monitors between February 22 and November 1, 2002 while a more in depth water quality survey of the Lake Wentworth deep sampling stations (Sites 1 Fullers, 2 Triggs and 12 Governors Deep) was conducted by the **Center for Freshwater Biology (CFB)** on August 1, 2002 to augment the volunteer monitoring data. Generally speaking, the year 2002 Lake Wentworth water quality remained high as characterized by the high water clarity, the relatively low levels of microscopic plant "algal" growth and the low phosphorus, nutrient, levels (Table 5).

For the second consecutive year, many participating New Hampshire Lakes Lay Monitoring Program lakes (including Lake Wentworth) were characterized by near record/record high water transparency measurements (Figures 23, 25 & 27). Furthermore, the 2002 microscopic plant "algal" levels included some of the lower concentrations documented over the past ten years of water quality monitoring for many participating lakes. The dry summer months limited the flushing of sediments and nutrients into our lakes; the same nutrients that make our lawns green will also stimulate plant growth in our New Hampshire lakes. Thus, the low levels of microscopic plant growth and the very clear waters that characterized most New Hampshire lakes are associated (at least in part) with the limited summer 2002 rainfall.

The following section discusses the year 2002 and historical Lake Wentworth water quality data. *Refer to Appendix A and Appendix B for a complete listing of the year 2002 Lake Wentworth water quality data.*

**Table 5: 2002 Lake Wentworth Seasonal Average Water Quality Readings and Water Quality Classification Criteria used by the New Hampshire Lakes Lay Monitoring Program.**

Parameter	Oligotrophic "Pristine"	Mesotrophic "Transitional"	Eutrophic "Enriched"	Lake Wentworth Average (range)	Lake Wentworth Classification
Water Clarity (meters)	> 4.0	2.5 - 4.0	< 2.5	7.5 meters (range: 6.4 - 9.5)	Oligotrophic
Chlorophyll a (ppb)	< 3.0	3.0 - 7.0	> 7.0	2.0 ppb (range: 0.7 - 5.3)	Oligotrophic
Phosphorus (ppb)	< 15.0	15.0 - 25.0	> 25.0	* 5.8 ppb (range: 5.7 - 6.0)	Oligotrophic

\* Total Phosphorus data were collected in the surface waters (epilimnion) on August 1, 2002.

**4) Water Clarity (measured as Secchi Disk transparency) –** The year 2002 Lake Wentworth water clarity values were high and generally remained well above the transparency of 13.2 feet (4 meters) that is considered the boundary between an unproductive "pristine" and more nutrient enriched "transitional" New Hampshire lake (Table 6 and Figures 15 - 20).

**Table 6: 2002 Water Clarity data summary for the Lake Wentworth deep sampling stations.**

Site	Seasonal Average Water Transparency (meters)
1 Fullers	7.8 meters (range: 6.5 - 9.5)
2 Triggs	7.5 meters (range: 6.5 - 8.2)
12 Governors	7.3 meters (range: 6.4 - 8.7)

A Comparison between the year 2002 and the historical (1984-2001) Lake Wentworth water transparency data indicate the seasonal average Secchi Disk transparency values were some of the higher values on documented (i.e. clearer water) at each of the three deep sampling stations (Figures 23, 25 and 27).

**5) Microscopic plant abundance “greenness” (measured as chlorophyll  $a$ )** – The year 2002 seasonal chlorophyll  $a$  values generally remained below the concentration of 3 parts per billion (ppb) that is considered the boundary between a nutrient poor and more nutrient enriched “greener” lake (Figures 15, 17 and 19). The 2002 seasonal average chlorophyll  $a$  concentrations were similar among the three Lake Wentworth deep sampling stations (Table 7).

**Table 7: 2002 Chlorophyll  $a$  data summary for the Lake Wentworth deep sampling stations.**

Site	Seasonal Average Chlorophyll $a$ (ppb)
1 Fullers	2.1 ppb (range: 1.3 – 5.3)
2 Triggs	1.9 ppb (range: 0.7 – 3.0)
12 Governors	2.0 ppb (range: 1.0 – 2.6)

The year 2002 seasonal chlorophyll  $a$  concentrations documented in Lake Wentworth, Site 1 Fullers, decreased relative to the 2001 seasonal average chlorophyll  $a$  concentration (Figure 24). On the other hand, the seasonal average chlorophyll  $a$  concentrations increased at Sites 2 Triggs and 12 Governors Deep, relative to the 2001 seasonal average concentrations (Figures 26 & 28). No new maximum or minimum chlorophyll  $a$  concentrations were documented in Lake Wentworth during the 2002 sampling season.

**6) Background (dissolved) water color: often perceived as a “tea” color in more highly stained lakes** – The year 2002 Lake Wentworth dissolved color concentration averaged 11.5 chloroplatinate units (cpu) and fell within the classification of an uncolored, no “tea stain”, lake (Table 8). Dissolved color, or true color as it is sometimes called, is indicative of dissolved organic carbon levels in the water (a by-product of microbial decomposition). Small increases in water color from the natural breakdown of plant materials in and around a lake are not considered to be detrimental to water quality. However, increased color can lower water transparency, and hence, change the public perception of water quality.

**Table 8. Dissolved Color Classification Criteria used by the New Hampshire Lakes Lay Monitoring Program.**

Range	Classification
0 - 10	Clear
10 - 20	Slightly colored
20 - 40	light tea color
40 - 80	tea colored
> 80	highly tea colored

**6) Total Phosphorus: the nutrient considered most responsible for elevated microscopic plant growth in our New Hampshire Lakes.** - Total phosphorus concentrations, measured in the surface waters (epilimnion), were low when measured in Lake Wentworth on August 1, 2002. The Lake Wentworth total phosphorus concentrations ranged from 5.7 to 6.0 parts per billion

(ppb) in the surface waters and remained well below the concentration of 10 ppb that is considered sufficient to cause an algal bloom.

Supplemental total phosphorus samples were collected at selected tributary sampling locations. The tributary total phosphorus concentrations were highly variable among streams and ranged from 5.0 to 23.2 parts per billion (ppb). *Refer to Appendix B for a complete listing of the 2002 Lake Wentworth total phosphorus data.* While elevated phosphorus concentrations can occur naturally, they can also be associated with improper land use activities (e.g. excessive fertilizer applications, construction without the proper erosion control measures in place) that result in “excessive” nutrient runoff into our lakes. Remember that the same nutrients that make our lawns green will also cause “greening” in our lakes. If you suspect problems around the ponds, we can institute a more thorough sampling program at selected points to help discern whether or not problems exist (contact Bob Craycraft @ 862-3696 for further information).

**7) Resistance against acid precipitation (measured as total alkalinity)** – The year 2002 Lake Wentworth alkalinity measured 6.3 milligrams per liter (mg/l) which is characteristic of a lake with a moderate vulnerability to acid precipitation according to the standards devised by the New Hampshire Department of Environmental Services (Table 9). Generally speaking, the geology of the region does not contain the mineral content (e.g. limestone) which increases the buffering capacity in our surface waters. Thus, lakes in the vicinity (i.e. Lake Winnepesaukee and Merrymeeting Lake) have naturally low alkalinities.

**Table 9. Alkalinity Classification  
Criteria used by the New Hampshire  
Department of Environmental  
Services**

Range	Classification
< 0	Acidified
0 -2	Extremely Vulnerable
2.1 - 10.0	Moderately Vulnerable
10.1 - 25.0	Low Vulnerability
> 25.0	Not Vulnerable

**Lake acidity (measured as pH)** - The year 2002 Lake Wentworth pH data, measured by the **Center for Freshwater Biology** on August 1, 2002, ranged from 6.8 to 7.2 units in the surface waters and remained well within the tolerable range for most aquatic organisms.

**8) Dissolved salts: measured as specific conductivity** – Specific Conductivity levels, documented in Lake Wentworth, were low and ranged from 54.0 to 63.0 micro-Siemans ( $\mu$ S) when measured at the deep, open water, sampling stations. High specific conductivity values can be an indication of problem areas around a lake where failing septic systems, heavy fertilizer applications and sedimentation contribute “excessive” nutrients that make their way into Lake Wentworth. High specific conductivity values can also be associated with road salt runoff that is flushed into our New Hampshire lakes.

**9) Temperature and dissolved oxygen profiles** – Temperature profiles collected by the volunteer monitors indicate Lake Wentworth becomes stratified

into three distinct thermal layers during the summer months; a warm upper water layer, the **epilimnion**, which overlies a layer of rapidly decreasing temperatures, the **thermocline**, and a deep cold-water layer, the **hypolimnion**. The formation of thermal stratification limits the replenishment of oxygen in the deeper waters and under adverse conditions can result in oxygen depletion near the lake-bottom.

**Dissolved oxygen concentrations required for a healthy fishery –**

Dissolved oxygen concentrations documented by the **Center for Freshwater Biology** and volunteer monitors were significantly reduced near the lakebottom during the period of summer temperature stratification (Appendix A and Table 10). Dissolved oxygen concentrations below 5 milligrams per liter are commonly considered inadequate for the successful growth and reproduction of coldwater fish species such as trout and salmon. The lake of dissolved oxygen in the deep, cold waters indicate Lake Wentworth is best suited as a warmwater fishery.

**Table 10. 2002 Lake Wentworth Dissolved Oxygen (DO) Concentrations and corresponding water quality classification criteria.**

Sampling Station & Date	DO Range (ppm) *	Classification
1 Fullers 09/08/02	0.0 – 1.5 ppm	“enriched”
2 Triggs 08/01/02	1.9 – 3.9 ppm	“transitional”
12 Governors 09/08/02	0.0 – 3.0 ppm	“enriched”

*\* Classification based on Dissolved oxygen Concentrations in the bottom waters (hypolimnion). Dissolved oxygen concentrations > 5 ppm are often considered typical of a “pristine” lake while dissolved oxygen concentrations < 2.0 ppm are considered typical of an “enriched” lake. Dissolved oxygen concentrations between 2.0 and 5.0 ppm are considered typical of a moderately productive “transitional” lake.*

8) Comparisons between the **Center for Freshwater Biology Group** and lay monitor data indicate the Lake Wentworth volunteer monitors are doing an excellent job of collecting water quality data.

9) Based on the current and historical water quality data, Lake Wentworth would be considered a water body that is transitioning between an unproductive “pristine” and a moderately nutrient enriched “transitional” New Hampshire lake. While the water clarity measurements and the microscopic plant “algal” concentrations have generally fallen within the ranges considered typical of a “pristine” lake, short term algal blooms, higher total phosphorus concentrations near the lakebottom and low dissolved oxygen concentrations near the lakebottom are characteristic of a moderately nutrient enriched, **mesotrophic**, lake. A first step towards preventing future water quality degradation in Lake Wentworth is to take action at the local level and do your part to minimize the number of pollutants (particularly sediment and the nutrient phosphorus) that enter the lake. Whenever possible, **maintain riparian buffers** (vegetative buffers adjacent to the water body). These buffers will biologically “take up” nutrients before the nutrients enter the lake and will also provide physical filters that allow materials to settle out before reaching the lake. **Reduce fertilizer applications.** Most residents apply far more fertilizers than necessary which can be a costly expense to the homeowner and can also be detrimental to the lake since the same nutrients that make our lawns green will also stimulate plant growth

in our lakes. **Make sure your septic system is well maintained.** Have your septic system pumped out on a regular basis. An improperly functioning septic system can contribute "excessive" nutrients into the lake and result in early failure, costing thousands of dollars to repair or replace. Future volunteer monitoring efforts should be directed at pinpointing problematic regions around the lake where corrective and educational efforts should be focused. It is important to make sure the watershed residents are well-educated on water quality related issues. Numerous publications are available through University of New Hampshire Cooperative Extension, the New Hampshire Lakes Association, the New Hampshire Department of Environmental Services as well as several other local, state and federal agencies. It is imperative that future activities within the Lake Wentworth watershed are carefully thought out before implementation if water quality degradation is to be minimized.

# COMMENTS AND RECOMMENDATIONS

---

1) We recommend that each participating association, including the Lake Wentworth Association, continue to develop its database on lake water quality through continuation of the long-term monitoring program. The database currently provides information on the short-term and long-term cyclic variability that occurs in the lake and through continued monitoring would enable more reliable predictions of both short-term and long-term water quality trends.

2) We recommend initiating lake sampling early in the season (April/May) to document Lake Wentworth's reaction to the nutrient and acid loadings that typically occur during and after spring thaw. Sampling should include alkalinity, chlorophyll *a*, dissolved color and Secchi Disk transparency measurements. Phosphorus samples are also recommended from both the in-lake and the tributary sampling sites. When tributary samples are collected, streamflow measurements should be included whenever possible.

3) Frequent "weekly" water quality samples, necessary to assess the current condition of Lake Wentworth, should continue to be collected whenever possible. Weekly sampling of chlorophyll *a*, Secchi Disk transparency, dissolved color and alkalinity is recommended while the addition of monthly total phosphorus samples would also be useful to track variations in nutrient loading during the summer months.

4) Follow up on the recommendations highlighted in the Lake Wentworth Diagnostic Study written and available through the New Hampshire Department of Environmental Services. The Lake Wentworth Diagnostic study targets certain locations within the watershed where corrective efforts should be targeted and also provides educational materials that can be distributed among local residents. When educational materials are to be distributed, focus on a watershed wide audience since all activities within the Lake Wentworth watershed will ultimately impact the lake.

# TABLE OF CONTENTS

---

PREFACE.....	I
ACKNOWLEDGMENTS .....	II
CRESCENT LAKE - 2002 NON-TECHNICAL SUMMARY .....	IV
LAKE WENTWORTH - 2002 NON-TECHNICAL SUMMARY .....	VIII
COMMENTS AND RECOMMENDATIONS.....	XIII
TABLE OF CONTENTS .....	XIV
REPORT FIGURES .....	XVI
INTRODUCTION.....	1
The New Hampshire Lakes Lay Monitoring Program.....	1
Importance of Long-term Monitoring .....	3
Purpose and Scope of This Effort .....	5
CLIMATIC SUMMARY - 2002 .....	6
Water Quality and the Weather .....	6
Precipitation (2001 and 2002) .....	7
Temperature (2002) .....	9
Water Quality Impacts .....	10
Water Transparency and Dissolved “tea” Colored Water .....	10
Sediment Loading.....	10
Nutrient Loading .....	11
Microscopic “Algal” and Macroscopic “Weed” Plant Growth.....	11
DISCUSSION OF LAKE AND STREAM MONITORING MEASUREMENTS.....	13
Thermal Stratification in the Deep Water Sites .....	13
Water Transparency .....	13
Chlorophyll <i>a</i> .....	14
Turbidity * .....	14
Dissolved Color.....	14
Total Phosphorus.....	15
Streamflow .....	15
pH *.....	15
Alkalinity.....	16
Specific Conductivity * .....	16
Dissolved Oxygen and Free Carbon Dioxide * .....	16
Underwater Light * .....	17
Indicator Bacteria * .....	17
Phytoplankton * .....	18
Zooplankton * .....	18
Macroinvertebrates * .....	19
Fish Condition .....	19
Zebra Mussels.....	20
UNDERSTANDING LAKE AGING .....	22



<b>(EUTROPHICATION).....</b>	<b>22</b>
How can you minimize your water quality impacts?.....	24
<b>BIOTOXINS: TOXIC CHEMICALS PRODUCED BY CYANOBACTERIA.....</b>	<b>26</b>
<b>REFERENCES.....</b>	<b>28</b>
<b>REPORT FIGURES .....</b>	<b>32</b>
<b>APPENDIX A .....</b>	<b>A-1</b>
<b>APPENDIX B .....</b>	<b>B-1</b>
<b>APPENDIX C .....</b>	<b>C-1</b>

# REPORT FIGURES

---

<b>Figure 1.</b> LLMP Objectives .....	1
<b>Figure 2.</b> Awards & Recognition .....	1
<b>Figure 3.</b> National LLMP Support Volunteer Monitoring Programs .....	2
<b>Figure 4.</b> Algal Standing Crop: 1988-1992.....	3
<b>Figure 5.</b> Algal Standing Crop: 1986-1995.....	4
<b>Figure 6.</b> Monthly Precipitation (1979-2002).....	7
<b>Figure 7.</b> Monthly Precipitation (1979-2002).....	8
<b>Figure 8.</b> Monthly Snowfall (1982-2002).....	8
<b>Figure 9.</b> Monthly Temperature (1984-2002).....	9
<b>Figure 10.</b> Typical Temperature Conditions: Summer .....	13
<b>Figure 11.</b> Location of the year 2002 Crescent Lake deep sampling station, Sites 6 Center, Wolfeboro New Hampshire.....	32
<b>Figure 12.</b> Location of the year 2002 Lake Wentworth deep sampling stations, Sites 1 Fullers, 2 Triggs, and 12 Governors Deep, Wolfeboro New Hampshire.....	34
<b>Figure 13.</b> Crescent Lake, 2002. Seasonal Secchi Disk (water transparency) and chlorophyll <i>a</i> trends for Site 6 Center.....	36
<b>Figure 14.</b> Crescent Lake, 2002. Seasonal Secchi Disk (water transparency) and dissolved color trends for Site 6 Center. ....	36
<b>Figure 15.</b> Lake Wentworth, 2002. Seasonal Secchi Disk (water transparency) and chlorophyll <i>a</i> trends for Site 1 Fullers.....	38
<b>Figure 16.</b> Lake Wentworth, 2002. Seasonal Secchi Disk (water transparency) and dissolved color trends for Site 1 Fullers.....	38
<b>Figure 17.</b> Lake Wentworth, 2002. Seasonal Secchi Disk (water transparency) and chlorophyll <i>a</i> trends for Site 2 Triggs. ....	40
<b>Figure 18.</b> Lake Wentworth, 2002. Seasonal Secchi Disk (water transparency) and dissolved color trends for Site 2 Triggs.....	40
<b>Figure 19.</b> Lake Wentworth, 2002. Seasonal Secchi Disk (water transparency) and chlorophyll <i>a</i> trends for Site 12 Governors Deep.....	42
<b>Figure 20.</b> Lake Wentworth, 2002. Seasonal Secchi Disk (water transparency) and dissolved color trends for Site 12 Governors Deep.....	42
<b>Figure 21.</b> Comparison of the year 2002 Crescent Lake, Site 6 Center, lay monitor Secchi Disk transparency data with historical water quality data collected in conjunction with the New Hampshire Lakes Lay Monitoring Program (1984-2001). ....	44

<b>Figure 22.</b> Comparison of the year 2002 Crescent Lake, Site 6 Center, lay monitor chlorophyll <i>a</i> data with historical water quality data collected in conjunction with the New Hampshire Lakes Lay Monitoring Program (1984-2001). .....	44
<b>Figure 23.</b> Comparison of the year 2002 Lake Wentworth, Site 1 Fullers, lay monitor Secchi Disk transparency data with historical water quality data collected in conjunction with the New Hampshire Lakes Lay Monitoring Program (1984-2001). .....	46
<b>Figure 24.</b> Comparison of the year 2002 Lake Wentworth, Site 1 Fullers, lay monitor chlorophyll <i>a</i> data with historical water quality data collected in conjunction with the New Hampshire Lakes Lay Monitoring Program (1984-2001). .....	46
<b>Figure 25.</b> Comparison of the year 2002 Lake Wentworth, Site 2 Triggs, lay monitor Secchi Disk transparency data with historical water quality data collected in conjunction with the New Hampshire Lakes Lay Monitoring Program (1984-2001). .....	48
<b>Figure 26.</b> Comparison of the year 2002 Lake Wentworth, Site 2 Triggs, lay monitor chlorophyll <i>a</i> data with historical water quality data collected in conjunction with the New Hampshire Lakes Lay Monitoring Program (1984-2001). .....	48
<b>Figure 27.</b> Comparison of the year 2002 Lake Wentworth, Site 12 Governors Deep, lay monitor Secchi Disk transparency data with historical water quality data collected in conjunction with the New Hampshire Lakes Lay Monitoring Program (1987-2001). ....	50
<b>Figure 28.</b> Comparison of the year 2002 Lake Wentworth, Site 12 Governors Deep, lay monitor chlorophyll <i>a</i> data with historical water quality data collected in conjunction with the New Hampshire Lakes Lay Monitoring Program (1987-2001). .....	50
<b>Figure 29.</b> Temperature and dissolved profiles collected in Crescent Lake, Site 6 Center, on August 1, 2002.....	52



# INTRODUCTION

## The New Hampshire Lakes Lay Monitoring Program

The 2002 sampling season marked the twenty-fourth anniversary for the NH Lakes Lay Monitoring Program (LLMP). The LLMP has grown from a university class project on Chocorua Lake and pilot study on the Squam Lakes to a comprehensive state-wide program with over 500 volunteer monitors and more than 100 lakes participating. Originally developed to establish a database for determining long-term trends of lake water quality for science and management, the program has expanded by taking advantage of the many resources that citizen monitors can provide (Figure 1).

The NH LLMP has gained an international reputation as a successful cooperative monitoring, education and research program. Current projects include: the use of volunteer generated data for non-point pollution studies using high tech analysis system (Geographic Information Systems and Satellite Remote Sensing), and intensive watershed monitoring for the development of watershed nutrient budgets, investigations of water quality and indicator organisms (food web analysis, fish condition, and stream invertebrates). The key ingredients responsible for the success of the program include innovative cost share funding and cost reduction, assurance of credible data, practical sampling protocols and, most importantly, the interest and motivation of our volunteer monitors.

The 2002 sampling season was another exciting year for the New Hampshire Lakes Lay Monitoring Program. National recognition for the high quality of work by you, the volunteer monitors, continued with awards, requests for pro-

Figure 1. LLMP Objectives




Figure 2. Awards & Recognition

- 1983 - NH Environmental Law Council Award
- 1984 - Governor's Volunteerism Award
- 1985 - CNN Science & Technology Today
- 1988 - Governor's "Gift" request funded
- 1990 - New Hampshire Journal TV Spot
- 1991 - Renew America Success Award
- Environmental Success Index
- UN Environmental Programme Award
- Soviet Embassy Reception
- White House Environment Briefing
- 1992 - EPA Administrators Award
- Environmental Exchange Network Listing
- 1993 - NH Lakes Association Award
- 1994 - Fourth National Citizens' Volunteer Monitoring Conference
- EPA Office of Watersheds Award
- 1995 - Co-editors Volunteer Monitor Newsletter (Spring Issue)
- Winnepesaukee Watershed Project
- 1996 - Fifth National Citizens' Volunteer Monitoring Conference
- 1998 - Governor's Proclamation for Anniversary
- Our New England Lakes Conference Host
- 1999 - EPA Watershed Academy Host
- 2000 - Sixth National Citizen's Monitoring Conference
- 2001 - Lake Chocorua Project Highlighted at National Lakes Conference and National Water Monitoring Conference



gram information and invitations to speak at national conferences (Figure 2).

Watersheds and water resources continue to be important topics covered in our spring semester Multidisciplinary Lakes Management course and our summer Community Mapping with GIS and Watershed Ecology courses that are held annually. These courses for educators, community leaders and other interested persons teach participants how to use the desktop Geographic Information System software - a computer based technology that allows users to create, map, manage, analyze and manipulate data and related locational information.

Research initiated by the New Hampshire Lakes Lay Monitoring Program and collaborators Dr. John Sasner and Dr. Jim Haney of the UNH Center for Freshwater Biology continued to focus on how watershed development and our activities on the landscape play a role in creating potentially toxic algae blooms. Analogous to the 'red tide' of estuaries, certain blue-green algae (microscopic bacteria) can produce toxins that are health risks to animals and humans. A \$150,000 grant from the Environmental Protection Agency and additional assistance from the University's Water Resources Research Center


has supported field trips to over 50 lakes throughout the state to monitor these populations, test for toxicity, trace the movement of toxins into mussels, crayfish and fish and determine what conditions favor toxic blooms of these organisms. Lakes Lay Monitoring Program participants helped make this EPA funded project a success by providing in-lake transportation and collecting water and mussel samples that were incorporated into ongoing research projects.

A United States Geological Survey Grant, administered through the UNH Water Resource Research Center, provided support for a review of historical nutrient/water budgets and the formulation of statewide phosphorus export coefficients for our New Hampshire Lakes. This five year grant focuses on how our activities on the land impact water quality. The recently completed volunteer facilitated water/nutrient budgets for Goose Pond and the Squam Lakes, as well as, other volunteer generated data, have been an integral part of the project's success.

A volunteer facilitated water quality improvement study has indicated significant reductions in phosphorus loadings due to berms, swales and diversions that were installed adjacent to Chocorua Lake to mitigate water quality problems. This project has received national recognition due to the cooperation among numerous agencies that include the NH Department of Transportation, the Natural Resources Conservation

**Figure 3. National LLMP Support to Volunteer Monitoring Programs**

**NH LLMP Directly involved with the Initiation, Expansion or Support of Volunteer Programs in 24 States.**



Light gray shading denotes LLMP assisted states


Service, North Country Resource and Development, the Carrol County Conservation District and the NH Department of Environmental Services. The new phase of this project includes further documentation of these best management practices to deal with road drainages as well as a new investigation that is determining the water quality protection provided by the wetland areas in the watershed.

We continue to be listed as a model citizen-monitoring program on the Environmental Success Index of Renew America, the Environmental Network Clearinghouse and the National Awards Council for Environmental Sustainability. To date, the approach and methods of the NH LLMP have been adopted by new or existing programs in twenty-four states and eleven countries (Figure 3)!


## Importance of Long-term Monitoring

A major goal of our monitoring program is to identify any short or long-term changes in the water quality of the lake. Of major concern is the detection of cultural eutrophication: increases in the productivity of the lake, the amount of algae and plant growth, due to the addition of nutrients from human activities. Changes in the natural buffering capacity of the lakes in the program is also a topic of great concern, as New Hampshire receives large amounts of acid precipitation, yet most of our lakes contain little mineral content to neutralize this type of pollution.

For over two decades, weekly data collected from lakes participating in the **New Hampshire Lakes Lay Monitoring Program** have indicated there is quite a variation in water quality indicators through the open water season (April through November) on the majority of lakes. Short-term differences may be due to variations in weather, lake use, or other chance events. Monthly sampling of a lake during a single summer provides some useful information, but there is a greater chance that important short-term events such as algal blooms or the lake's response to storm run-off will be missed. These short-term fluctuations may be unrelated to the actual long-term trend of a lake or they may be indicative of the changing status or "health" of a lake.



Consider the hypothetical data depicted in Figure 4. Sampling only once a year during August from 1988 to 1992 produced a plot suggesting a decrease in eutrophication. However, the actual long-term term trend of the lake, increasing eutrophy, can only be clearly discerned by frequent sampling over a ten year period (Figure 5). In this instance, the information necessary to distinguish between short-term fluctuations "noise" and long-term trends "signal" could only be accomplished through the frequent collection of water quality data over many years. To that end, the establishment of a long term database was essential to trend detection.



The number of seasons it takes to distinguish between the noise and the signal is not the same for each lake. Evaluation and interpretation of a long-term data base will indicate that the water quality of the lake has worsened, improved, or remained the same. In addition, different areas of a lake may show a different response. As more data are collected, prediction of current and future trends can be made. No matter what the outcome, this information is essential for the intelligent management of your lake.

There are also short-term uses for lay monitoring data. The examination of different stations in a lake can disclose the location of specific problems and corrective action can be initiated to handle the situation before it becomes more serious. On a lighter note, some associations post their weekly data for use in determining the best depths for finding fish!

It takes a considerable amount of effort as well as a deep concern for one's lake to be a volunteer in the **NH Lakes Lay Monitoring Program**. Many times a monitor has to brave inclement weather or heavy boat traffic to collect samples. Sometimes it even may seem that one week's data is just the same as the next week's data. Yet every sampling provides important information on the variability of the lake.

We are pleased with the interest and commitment of our Lay Monitors and are proud that their work is what makes the **NH LLMP** the most extensive, and we believe, the best volunteer program of its kind.



## Purpose and Scope of This Effort

The year 2002 was the nineteenth year that water quality monitoring of Lake Wentworth and Crescent Lake was undertaken in conjunction with the **NH Lakes Lay Monitoring Program (NH LLMP)**. The monitoring program is designed to establish a long-term database to which future data could be compared and to identify localized problem areas where future educational and corrective efforts should be focused. Sampling emphasis was placed on deep open water sampling stations in both Lake Wentworth and Crescent Lake while additional sampling of selected tributary inlets was also conducted during the 2002 sampling season.

The primary purpose of annual lake reporting is to discuss results of the current monitoring season with emphasis on current conditions of New Hampshire lakes including the extent of eutrophication and the lakes' susceptibility to increasing acid precipitation. This information is part of a large data base of historical and more recent data compiled and entered onto computer files for New Hampshire lakes that include New Hampshire Fish and Game surveys of the 1930's, the surveys conducted by the New Hampshire Water Supply and Pollution Control Commission and the **CFB/FBG** surveys. However, care must be taken when comparing current results with early studies. Many complications arise due to methodological differences of the various analytical facilities and technological improvements in testing.

# Climatic Summary - 2002

---

## Water Quality and the Weather

Since the inception of the **New Hampshire Lakes Lay Monitoring Program (NH LLMP)**, questions have been asked pertaining to water quality changes that occur in our New Hampshire lakes and ponds. The most commonly noticed changes are those associated with decreasing water clarities, increasing algal growth (greenness), and increasing plant growth around the lake's periphery. Over the long haul, changes such as these are attributed to a lake's natural aging process; what is known as "**eutrophication**". However, short-term water quality changes such as those mentioned above are often encountered even in our most pristine lakes and ponds. These water quality changes often coincide with variations in weather variables such as precipitation and temperature.

Climatic "swings" can have a profound effect on water quality, sometimes positive and other times negative. For instance, 1996 was a wet year relative to other years of **LLMP** water quality monitoring. This translated into reduced water clarities, elevated microscopic plant "algal" growth and increased total phosphorus concentrations for most participating **LLMP** lakes. Past monitoring through the **NH LLMP** has recognized that wet years such as 1996 often result in poorer water quality in many of our New Hampshire lakes, relative to years with more typical precipitation levels. "Excessive" runoff associated with wet periods often facilitates the transport of pollutants such as nutrients (including phosphorus), sediment, dissolved colored compounds, as well as toxic materials such as herbicides, automotive oils, etc. into water bodies. As a result, lakes often respond with shallower (less clear) water clarities and elevated algal abundance "greenness" during these periods. Similarly, short-term storm events can have a profound effect on the water quality. Take for instance the "100 year storm" (October 21-22, 1996) that blanketed southern New Hampshire with approximately 6 inches of rain over a 30-hour period. This storm resulted in increased sedimentation and organic matter loading into our lakes as materials were flushed into the water bodies from the adjacent landscape. Likewise, the heavy rains that saturated the soil and resulted in flood conditions in June 1998 (heaviest rains occurring on June 12 and 13) resulted in significantly shallower water transparency readings in the weeks to months that followed. While events such as the October 1996 and the June 1998 storms are short lived, they can have a profound effect on our water quality in the weeks to months that follow, particularly when nutrients that stimulate plant growth are retained in the lake.


**NH LLMP** data collected during dry years such as 1985, on the other hand, coincide with improved water quality for many New Hampshire lakes. Reduced transport of pollutants into the lake often results in higher water quality measured as deeper water transparencies, lower microscopic plant "algae" concentrations and lower nutrient concentrations. Do all lakes experience poorer water quality as a result of heavy precipitation events? Simply stated, the answer is no. While most New Hampshire lakes are characterized by reduced water clarities, increased nutrients and elevated plant "algal" concentrations following periods, or years, of heavy precipitation, a handful of lakes actually benefit from these types of events. These are generally lakes characterized by high nutrient concentrations and high "algal" concentrations that are diluted by water-

shed runoff and thus benefit during periods, or years, of heavy rainfall. However, these more nutrient enriched lakes might be susceptible to nutrients entering the lake from seepage sources such as poorly functioning septic systems.


### Precipitation (2001 and 2002)

The total 2001 precipitation (measured as water equivalent) measured 32.82 inches and was over eight inches below the long-term precipitation average of 41.06 inches that was documented over the past 24 years: 1979-2002. Furthermore, the monthly precipitation documented between April and December 2001 was generally well below average and resulted in a low groundwater table and drought-like conditions from the summer into the fall months (Figure 6). Monthly rainfall was slightly above average between February and June, 2002 while the rainfall was significantly below normal levels during the months of July and August (Figure 7). The dry conditions that began during the summer of 2001 resulted in a low water table that continued into the fall of 2002 and translated into minimal overland runoff during much of the 2002 sampling season.


Snowfall also had an impact on the minimal overland runoff during the 2002 sampling season. The 2002 snowfall during the months of January, February, March and April were significantly below the 21-year average: 1982-2002 (Figure 8). Significant accumulations of winter snowpack can result in a period of heavy overland runoff in the spring that oftentimes coincides with significant sediment and nutrient loading that negatively impact water quality. The lack of significant spring snowpack in 2002 reduced the spring sediment and nutrient load to many of our lakes and streams.



**Figure 7: Lakeport 2 Climatological  
Sampling Station (Laconia, NH)  
Monthly Precipitation (1979-2002)**



**Figure 8: Lakeport 2 Climatological  
Sampling Station (Laconia, NH)  
Monthly Snowfall (1982-2002)**




### Temperature (2002)

Similar to the impact of precipitation extremes, temperature extremes can have far reaching effects on the water quality, particularly early in the year and during the summer months. Atypically warm spells can account for a rapid snowpack melt resulting in flooding and a massive influx of materials (e.g. nutrients, sediments) into our lakes during the late winter and early spring months. Early spring runoff periods coincide with minimal vegetative cover (that acts as a pollutant filter and soil stabilizer) and thus leaves the landscape highly susceptible to erosion. As we progress into the summer months, atypically warm periods can enhance both microscopic “algal” and macroscopic “aquatic weed” plant growth. During the summer growing season, above average temperatures often result in algal blooms that under optimal conditions can reach nuisance proportions. These can include surface algal “scums” that cover the lake and wash up on the windward lakeshores.

During years such as 1994 and 1995, when above average temperatures characterized the summer months, participating NH LLMP lakes were generally characterized by increased algal concentrations, particularly in the shallows, where filamentous cotton candy like clouds of algae (i.e. *Mougeotia*) flourished. Other NH LLMP lakes had increased algal growth “greenness” and shallower water transparencies during these “hot” periods.

The 2002 average monthly temperatures during January, February, March and April were above the nineteen-year average temperatures: 1984-2002 (Figure 9). The above average temperatures coincided with less winter/early spring snowfall than normal (much of the precipitation fell as rain) and minimized the accumulation of winter snowpack. The above average winter and early spring temperatures also translated into



an early ice-out for our New Hampshire lakes relative to most years. The 2002 average monthly temperatures documented during the months of May (54.0°F) and June (64.1°F) were slightly below average while the 2002 average monthly temperatures during the months of July (72.1°F), August (73.0°F) and September (65.8°F) exceeded the nineteen year temperature average by 1.2°F to 4.8°F. The combination of the warm air temperatures early in the year and the above average temperatures between July and September coincided with above average lakewater temperatures during much of the summer sampling period.

## **Water Quality Impacts**

### **Water Transparency and Dissolved “tea” Colored Water**

As previously indicated, shallower water transparency readings are characteristic of most New Hampshire lakes during wet years and following short term precipitation events. Wet periods often coincide with greater concentrations of dissolved “tea” colored compounds (dissolved organic matter resulting from the breakdown of vegetation and soils) washed in from surrounding forests and wetlands. Dissolved water color is not indicative of water quality problems (although large increases in dissolved color sometimes follow large land clearing operations) but in some of our more pristine program lakes, it nevertheless has a large effect on water clarity changes. Data collected by the **Center For Freshwater Biology (CFB)** since 1985 indicate most lakes are characterized by higher dissolved “tea” colored water during wet years relative to years more typical in terms of annual precipitation levels. In some of our more highly “tea” colored lakes the early spring months are also characterized by higher dissolved color concentrations, relative to mid-summer levels, due to the heavy runoff periods that flush highly colored water into our lakes during the period of spring snowmelt and following heavy spring rains.

### **Sediment Loading**

Sediments are continuously flushed into our lakes and ponds during periods of heavy watershed runoff, particularly during snowmelt and again during and following sporadic storm events during the summer and fall months. Many New Hampshire lakes experience water clarity decreases following storm events such as those described above. Lakes, ponds and rivers are particularly susceptible to sediment loadings in the early spring months when vegetated shoreside buffers, often referred to as riparian buffers, are reduced. With limited vegetation to trap sediments and suspended materials, a high percentage of the particulate debris and dissolved materials are flushed into the lake. Human activities such as logging, agriculture, construction and land clearing can also increase sediment displacement during and following heavy storm events throughout the year. These activities are often associated with excessive sediment loading in many of our lakes and ponds. As these materials (sediments) are transported into surface waters they can degrade water quality in a number of ways. When fine sediments (silt) enter a lake they tend to remain in the water column for relatively long periods of time. These suspended sediments can be abrasive to fish gills, ultimately leading to fish kills. Suspended sediments also reduce the available light necessary for plant growth that can result in plant die-offs and the subsequent oxygen depletion under extreme conditions.

As sediments settle out of the water column they can smother bottom dwelling aquatic organisms and fish spawning habitat. As the dead materials begin to decay the result can be noxious odors as well as stimulation of nuisance plant growth (i.e. scums along the lakebottom; new macroscopic plant growth). Note: one should keep in mind that nuisance plants such as water milfoil (*Myriophyllum heterophyllum*) will generally regenerate more rapidly than more favorable plant forms. This can result in more problematic weed beds than those present before the disturbance. Habitat changes associated with the accumulation of fine sediments and associated "muck" might also favor increased nuisance plant growth in the future. Another unfavorable attribute of sediment loading is that the sediments tend to carry with them other sorts of contaminant such as pathogens, nutrients and toxic chemicals (i.e. herbicides and pesticides).

Early symptoms of excessive sediment runoff include deposits of fine material along the lakebottom, particularly in close proximity to tributary inlets and disturbed regions previously discussed (i.e. construction sites, logging sites, etc.). Silt may be visible covering rocks or aquatic vegetation along the lakebottom. During periods of heavy overland runoff the water might appear brown and turbid which reflects the sediment load. As material collects along the lakebottom you might notice a change in the weed composition reflecting a change in the substrate type (note: aquatic plants will display natural changes in abundance and distribution, so be careful not to jump to hasty conclusions). If excessive sediment loading is suspected, take a closer look in these areas and assess whether or not the change is associated with sediment loading (look for the warning signs discussed above) or whether the changes might be attributable to other factors.

### **Nutrient Loading**

Nutrient loading is often greatest during heavy precipitation events, particularly during the periods of heavy watershed runoff. Phosphorus is generally considered the limiting nutrient for excessive plant and algal growth in New Hampshire lakes. Elevated phosphorus concentrations are generally most visible when documented in our tributary inlets where nutrients are concentrated in a relatively small volume of water. Much of the phosphorus entering our lakes is attached to particulate matter (i.e. sediments, vegetative debris), but may also include dissolved phosphorus associated with fertilizer applications and septic system discharge.

### **Microscopic "Algal" and Macroscopic "Weed" Plant Growth**

Historical Lakes Lay Monitoring Program data indicate most lakes experience "algal blooms" during years with above average summer temperatures (June, July and August) while years with heavy precipitation are also associated with an increased frequency and occurrence of "algal blooms" among participating LLMP lakes. Algal blooms are often green water events associated with decreases in water clarity due to their ability to absorb and scatter light within the water column, but can also accumulate near the lake bottom in shallow areas as "mats" or on the water surface as "scums" and "clouds". During some years, such as 1996, the algal blooms are predominantly green water events composed of algae distributed within the water column. New Hampshire lakes were particularly susceptible to algal blooms in 1996 as a function of the heavy runoff associated with the atypically wet year. Wet years such as 1996 can be particularly hard on lakes where excessive fertilizer applications, agricultural practices and construction activities favor the displacement of nutrients into surface waters. The

occasional formation of certain algal blooms is a naturally occurring phenomenon and is not necessarily associated with changes in lake productivity. However, increases in the occurrence of bloom conditions can be a sign of eutrophication (the "greening" of a lake). Shifts from benign (clean water) forms to nuisance (polluted water) cyanobacterial forms such as *Anabaena*, *Aphanizomenon* and *Oscillatoria*, can also be a warning sign that improper land use practices are contributing excessive nutrients into the lake.

Filamentous cotton-candy like "clouds" of the nuisance green algae, *Mougeotia* and related species, have been well documented in 1994 and 1995 when the temperatures during the months of June and July were well above normal. These algal "clouds" often develop within nearshore weed beds where they can be seen along the lakebottom and tend to flourish during warm periods. During cooler years, this type of algal growth is kept "in check" and generally does not reach nuisance proportions.

In other lakes, metalimnetic algae, algae which tend to grow in a thin layer along the thermocline gradient in a lake's middle depths, sometimes migrate up towards the lake surface causing a "bloom" event. If these algae are predominantly "nuisance" forms, like certain green or blue-green algae, they can be an early indication of nutrient loading.



# DISCUSSION OF LAKE AND STREAM MONITORING MEASUREMENTS

---

The section below details the important concepts involved for the various testing procedures used in the **New Hampshire Lakes Lay Monitoring Program**. Certain tests or sampling performed at the time of the optional **Center for Freshwater Biology** field trip are indicated by an asterisk (\*).


## Thermal Stratification in the Deep Water Sites

Lakes in New Hampshire display distinct patterns of temperature stratification, that develop as the summer months progress, where a layer of warmer water (the **epilimnion**) overlies a deeper layer of cold water (**hypolimnion**). The layer that separates the two regions characterized by a sharp drop in temperature with depth is called the **thermocline** or **metalimnion** (Figure 10). Some shallow lakes may be continually mixed by wind action and will never stratify. Other lakes may only contain a developed epilimnion and metalimnion.

## Water Transparency

Secchi Disk depth is a measure of the water transparency. The deeper the depth of Secchi Disk disappearance, the more transparent the lake water; light penetrates deeper if there is little dissolved and/or particulate matter (which includes both living and non-living particles) to absorb and scatter it.

In the shallow areas of many lakes, the Secchi Disk will hit bottom before it is able to disappear from view (what is referred to as a "Bottom Out" condition). Thus, Secchi Disk measurements are generally taken over the deepest sites of a lake. Transparency values greater than 4 meters are typical of clear, unproductive lakes while transparency values less than 2.5 meters are generally an indication of highly productive lakes. Water transparency values between 2.5 meters and 4 meters are generally considered indicative of moderately productive lakes.



## Chlorophyll *a*

The chlorophyll *a* concentration is a measurement of the standing crop of phytoplankton and is often used to classify lakes into categories of productivity called trophic states. **Eutrophic** lakes are highly productive with large concentrations of algae and aquatic plants due to nutrient enrichment. Characteristics include accumulated organic matter in the lake basin and lower dissolved oxygen in the bottom waters. Summer chlorophyll *a* concentrations average above 7 mg m<sup>3</sup> (7 milligrams per cubic meter; 7 parts per billion). **Oligotrophic** lakes have low productivity and low nutrient levels and average summer chlorophyll *a* concentrations that are generally less than 3 mg m<sup>3</sup>. These lakes generally have cleaner bottoms and high dissolved oxygen levels throughout. **Mesotrophic** lakes are intermediate in productivity with concentrations of chlorophyll *a* generally between 3 mg m<sup>3</sup> and 7 mg m<sup>3</sup>. Testing is sometimes done to check for **metalimnetic algal populations**, algae that layer out at the thermocline and generally go undetected if only epilimnetic (point or integrated) sampling is undertaken. Chlorophyll concentrations of a water sample collected in the thermocline is compared to the integrated epilimnetic sample. Greater chlorophyll levels of the point sample, in conjunction with microscopic examination of the samples (see Phytoplankton section below), confirm the presence of such a population of algae. These populations should be monitored as they may be an indication of increased nutrient loading into the lake.

## Turbidity \*

Turbidity is a measure of suspended material in the water column such as sediments and planktonic organisms. The greater the turbidity of a given water body the lower the Secchi Disk transparency and the greater the amount of particulate matter present. Turbidity is measured as nephelometric turbidity units (NTU), a standardized method among researchers. Turbidity levels are generally low in New Hampshire reflecting the pristine condition of the majority of our lakes and ponds. Increasing turbidity values can be an indication of increasing lake productivity or can reflect improper land use practices within the watershed which destabilize the surrounding landscape and allow sediment flushing into the lake.

While Secchi Disk measurements will integrate the clarity of the water column from the surface waters down to the depth of disappearance, turbidity measurements are collected at discrete depths from the surface down to the lake bottom. Such discrete sampling can identify layering algal populations (previously discussed) that are undetectable when measuring Secchi Disk transparency alone.

## Dissolved Color

The dissolved color of lakes is generally due to dissolved organic matter from **humic substances**, which are naturally-occurring polyphenolic compounds leached from decayed vegetation. Highly colored or "stained" lakes have a "tea" color. Such substances generally do not threaten water quality except as they diminish sunlight penetration into deep waters. Increases in dissolved watercolor can be an indication of increased development within the watershed as many land clearing activities (construction, deforestation, and the resulting increased run-off) add additional organic material to lakes. Natural fluctuations of dissolved color occur when storm events increase drainage from wetlands areas within the watershed. As suspended sediment is a diffi-

cult and expensive test to undertake, both dissolved color and chlorophyll information are important when interpreting the Secchi Disk transparency

Dissolved color is measured on a comparative scale that uses standard chloroplatinate dyes and is designated as a color unit or ptu. Lakes with color below 10 ptu are very clear, 10 to 20 ptu are slightly colored, 20 to 40 ptu are lightly tea colored, 40 to 80 ptu are tea colored and greater than 80 ptu indicates highly colored waters. Generally the majority of New Hampshire lakes have color between 20 to 30 ptu.

## **Total Phosphorus**

Of the two "nutrients" most important to the growth of aquatic plants, nitrogen and phosphorus, it is generally observed that phosphorus is the more limiting to plant growth, and therefore the more important to monitor and control. Phosphorus is generally present in lower concentrations, and its sources arise primarily through human related activity in a watershed. Nitrogen can be fixed from the atmosphere by many bloom-forming blue-green bacteria, and thus it is difficult to control. The total phosphorus includes all dissolved phosphorus as well as phosphorus contained in or adhered to suspended particulates such as sediment and plankton. As little as 10 parts per billion of phosphorus in a lake can cause an algal bloom.

Generally, in the more pristine lakes, phosphorus values are higher after spring melt when the lake receives the majority of runoff from its surrounding watershed. The nutrient is used by the algae and plants which in turn die and sink to the lake bottom causing surface water phosphorus concentrations to decrease as the summer progresses. Lakes with nutrient loading from human activities and sources (Agriculture, Logging, Sediment Erosion, Septic Systems, etc.) will show greater concentrations of nutrients as the summer progresses or after major storm events.

## **Streamflow**

Streamflow, when collected in conjunction with depth contour information, is a measure of the volume of water traversing a given stream stretch over a period of time and is often expressed as cubic meters per second. Knowledge of the streamflow is important when determining the amount of nutrients and other pollutants that enter a lake. Knowledge of the streamflow in conjunction with nutrient concentrations, for instance, will provide the information necessary to calculate phosphorus loading values and will in turn be useful in discerning the more impacted areas within a watershed.

## **pH \***

The pH is a way of expressing the acidic level of lake water, and is generally measured with an electrical probe sensitive to hydrogen ion activity. The pH scale has a range of 1 (very acidic) to 14 (very "basic" or alkaline) and is logarithmic (i.e.: changes in 1 pH unit reflect a ten times difference in hydrogen ion concentration). Most aquatic organisms tolerate a limited range of pH and most fish species require a pH of 5.5 or higher for successful growth and reproduction.

## Alkalinity

Alkalinity is a measure of the buffering capacity of the lake water. The higher the value the more acid that can be neutralized. Typically lakes in New Hampshire have low alkalinities due to the absence of carbonates and other natural buffering minerals in the bedrock and soils of lake watersheds.

Decreasing alkalinity over a period of a few years can have serious effects on the lake ecosystem. In a study on an experimental acidified lake in Canada by Schindler, gradual lowering of the pH from 6.8 to 5.0 in an 8-year period resulted in the disappearance of some aquatic species, an increase in nuisance species of algae and a decline in the condition and reproduction rate of fish. During the first year of Schindler's study the pH remained unchanged while the alkalinity declined to 20 percent of the pre-treatment value. The decline in alkalinity was sufficient to trigger the disappearance of zooplankton species, which in turn caused a decline in the "condition" of fish species that fed on the zooplankton.

The analysis of alkalinity employed by the **Center for Freshwater Biology** includes use of a dilute titrant allowing an order of magnitude greater sensitivity and precision than the standard method. Two endpoints are recorded during each analysis. The first endpoint (gray color of dye; pH endpoint of 5.1 ) approximates low level alkalinity values, while the second endpoint (pink dye color; pH endpoint of 4.6) approximates the alkalinity values recorded historically, such as NH Fish and Game data, with the methyl-orange endpoint method.

The average alkalinity of lakes throughout New Hampshire is low, approximately 6.5 mg per liter (calcium carbonate alkalinity). When alkalinity falls below 2 mg per liter the pH of waters can greatly fluctuate. Alkalinity levels are most critical in the spring when acid loadings from snowmelt and run-off are high, and many aquatic species are in their early, and most susceptible, stages of their life cycle.

## Specific Conductivity \*

The specific conductance of a water sample indicates concentrations of dissolved salts. Leaking septic systems and deicing salt runoff from highways can cause high conductivity values. Fertilizers and other pollutants can also increase the conductivity of the water. Conductivity is measured in micromhos (the opposite of the measurement of resistance ohms) per centimeter, more commonly referred to as micro-Siemans ( $\mu$ S).

## Dissolved Oxygen and Free Carbon Dioxide \*

Oxygen is an essential component for the survival of aquatic life. Submergent plants and algae take in carbon dioxide and create oxygen through **photosynthesis** by day. **Respiration** by both animals and plants uses up oxygen continually and creates **carbon dioxide**. Dissolved oxygen profiles determine the extent of declining oxygen concentrations in the lower waters. High carbon dioxide values are indicative of low oxygen conditions and accumulating organic matter. For both gases, as the temperature of the water decreases, more gas can be dissolved in the water.

The typical pattern of clear, unproductive lakes is a slight decline in hypolimnetic oxygen as the summer progresses. Oxygen in the lower waters is important for maintaining a fit, reproducing, cold water fishery. Trout and salmon generally require oxygen concentrations above 5 mg per liter (parts per million) in the cool deep waters.

On the other hand, carp and catfish can survive very low oxygen conditions. Oxygen above the lake bottom is important in limiting the release of nutrients from the sediments and minimizing the collection of undecomposed organic matter.

Bacteria, fungi and other **decomposers** in the bottom waters break down organic matter originating from the watershed or generated by the lake. This process uses up oxygen and produces carbon dioxide. In lakes where organic matter accumulation is high, oxygen depletion can occur. In highly stratified eutrophic lakes the entire hypolimnion can remain unoxygenated or **anaerobic** until fall mixing occurs.

The oxygen peaks occurring at surface and mid-lake depths during the day are quite common in many lakes. These characteristic **heterograde oxygen curves** are the result of the large amounts of oxygen, the by-product of photosynthesis, collecting in regions of high algal concentrations. If the peak occurs in the thermocline of the lake, metalimnetic algal populations (discussed above) may be present.

## Underwater Light \*

Underwater light available to photosynthetic organisms is measured with an **underwater photometer** which is much like the light meter of a camera (only waterproofed!). The **photic zone** of a lake is the volume of water capable of supporting photosynthesis. It is generally considered to be delineated by the water's surface and the depth that light is reduced to one percent surface irradiance by the absorption and scattering properties of the lake water. The one percent depth is sometimes termed the **compensation depth**. Knowledge of light penetration is important when considering lake productivity and in studies of submerged vegetation. Discontinuity (abrupt changes in the slope) of the profiles could be due to metalimnetic layering of algae or other particulates (discussed above). The underwater photometer allows the investigator to measure light at depths below the Secchi Disk depth to supplement the water clarity information.

## Indicator Bacteria \*

Certain disease causing organisms, pathogenic bacteria, viruses and parasites, can be spread through contact with polluted waters. Faulty septic systems, sewer leaks, combined sewer overflows and the illegal dumping of wastes from boats can contribute fecal material containing these pathogens. Typical water testing for pathogens involves the use of detecting coliform bacteria. These bacteria are not usually considered harmful themselves but they are relatively easy to detect and can be screened for quickly. Thus, they make good surrogates for the more difficult to detect pathogens.

**Total coliform** includes all coliform bacteria that arise from the gut of animals or from vegetative materials. **Fecal coliform** are those specific organisms that inhabit the gut of warm blooded animals. Another indicator organism **Fecal streptococcus** (sometimes referred to as **enterococcus**) also can be monitored. The ratio of fecal coliform to fecal strep may be useful in suggesting the type of animal source responsible for the contamination. In 1991, the State of New Hampshire changed the indicator organism of preference to E. Coli which is a specific type of fecal coliform bacteria thought to be a better indicator of human contamination. The new state standard requires Class A "bathing waters" to be under 88 organisms (referred to as colony forming units; cfu) per 100 milliliters of lakewater.

Ducks and geese are often a common cause of high coliform concentrations at specific lake sites. While waterfowl are important components to the natural and aesthetic qualities of lakes that we all enjoy, it is poor management practice to encourage these birds by feeding them. The lake and surrounding area provides enough healthy and natural food for the birds and feeding them stale bread or crackers does nothing more than import additional nutrients into the lake and allows for increased plant growth. As birds also are a host to the parasite that causes "swimmers itch", waterfowl roosting areas offer a greater chance for infestation to occur. Thus while leaving offerings for our feathered friends is enticing, the results can prove to be detrimental to the lake system and to human health.

## **Phytoplankton \***

The planktonic community includes microbial organisms that represent diverse life forms, containing photosynthetic as well as non-photosynthetic types, and including bacteria, algae, crustaceans and insect larvae (the insect larvae and zooplankton are discussed below in separate sections). Because planktonic algae or "phytoplankton" tend to undergo rapid seasonal cycles on a time scale of days and weeks, the levels of populations found should be considered to be most representative of the time of collection and not necessarily of other times during the ice-free season, especially the early spring and late fall periods.

The composition and concentration of phytoplankton can be indicative of the trophic status of a lake. Seasonal patterns do occur and must be considered. For example **diatoms**, tend to be most abundant in April-June and October-November, in the surface or epilimnetic layers of New Hampshire lakes. As the summer progresses, the dominant types might shift to **green algae** or **golden algae**. By late season **Blue-green bacteria** generally dominate. In nutrient rich lakes, nuisance green algae and/or bluegreen bacteria might dominate continually. After fall mixing diatoms might again be found to bloom.

## **Zooplankton \***

There are three groups of zooplankton that are generally prevalent in lakes: the **protozoa**, **rotifers** and **crustaceans**. Most research has been devoted to the last two groups although protozoa may be found in substantial amounts. Of the rotifers and the crustaceans, time and budgetary constraints usually make it necessary to sample only the larger zooplankton (macrozooplankton; larger than 80 or 150 microns; 1 million microns make up a meter). Thus, zooplankton analysis is generally restricted only to the larger crustaceans. Crustacean zooplankton are very sensitive to pollutants and are commonly used to indicate the presence of toxic substances in water. The crustaceans can be divided into two groups, the **cladocerans** (which include the "water fleas") and the **copepods**.

Macrozooplankton are an important component in the lake system. The filter feeding of the herbivorous ("grazing") species may control the population size of selected species of phytoplankton. The larger zooplankton can be an important food source for juvenile and adult planktivorous fish. All zooplankton play a part in the recycling of nutrients within the lake. Like the phytoplankton, zooplankton, tend to undergo rapid seasonal cycles. Thus, the zooplankton population density and diversity should be consid-

ered to be most representative of the time of collection and not necessarily of other times during the ice-free season, especially the early spring and late fall periods.

## Macroinvertebrates \*

Macroinvertebrates generally refer to the aquatic insect community living near the bottom substrate (i.e. sediments) while other invertebrate groups such as the crayfish, leeches and the aquatic worms are also included. Like the phytoplankton and zooplankton, previously discussed, the macroinvertebrates undergo seasonal cycles and are most representative of conditions for particular periods of the year. The mayflies are probably the most well known example of a seasonal aquatic macroinvertebrate as mayfly populations metamorphosize into adults as the water temperatures increase in the spring and thus giving rise to the name "mayflies". Macroinvertebrates are also sensitive to environmental conditions such as streamflow, temperature and food availability and are most representative of particular habitats along the stream continuum (i.e. some organisms prefer slower moving stream reaches while others prefer rapidly flowing waters).

Macroinvertebrates are an essential component to a healthy aquatic habitat. Macroinvertebrates help decompose organic matter entering the system such as leaves and twigs and also serve as a food source for many fish species.

While some macroinvertebrates are capable of breathing air as we do, others have gills and utilize oxygen dissolved in the water much as fish do. Macroinvertebrates also vary in their tolerance to depleting dissolved oxygen concentrations making them a good indicator of pollutants coming into the water body. The caddis flies (Trichoptera), the mayflies (Ephemeroptera) and the stoneflies (Plecoptera) are often considered highly sensitive to pollution while the "true" flies (Diptera) are often considered highly tolerant to pollution. However, exceptions to the above categorizations are often encountered.

A variety of indices have been proposed to characterize water bodies over a gradient of pollution levels ranging from least polluted to most polluted scenarios and often designated by assigning a numerical delineator (i.e. 1 is least polluted while 10 is most polluted). Such an index, the Hilsenhoff Biotic Index (HBI), or a modification thereof, is commonly used by stream monitoring programs around the country. Macroinvertebrate data are useful in discerning the more impacted areas within the watershed where corrective efforts should be directed. Unlike chemical measurements that represent ambient conditions in the water body, the macroinvertebrate community composition integrates the water quality conditions over a longer period (months to years) and can identify "hot" spots missed by chemical sampling. If you are interested in more information regarding macroinvertebrate monitoring contact the **LLMP** coordinator.

## Fish Condition

The assessment of fish species "health" is another biological indicator of water quality. Because fish are at the top of the food chain, their condition should reflect not only water quality changes that affect them directly but also those changes that affect their food supply. The fish condition index utilized by the **New Hampshire Fish Condition Program** is based on two components; fish scale analysis and a fish condition index.

Like tree trunks, fish scales have annual growth rings (annuli) that reflect their growth history and hence, provide a long-term record of past conditions in the lake. The fish condition index, based upon length and weight measurements, is a good indicator of the fish's health at the time of collection.

The resulting fish condition data can be compared among different lakes or among different years, or the index for a particular species can be compared to standard length-to-weight relationships that have been developed by fisheries biologists for many important fish species. In the end, the "health" of the various fish species reflects the overall water quality in the respective lake or pond.

## Zebra Mussels

Zebra mussels (*Dreissena polymorpha*) are non-native, freshwater mollusks. The veligers (larval form) are free swimming, nearly invisible, and profuse. Adult zebra mussel shells are elongate (D-shaped), about the size of a thumbnail and are usually striped. Zebra Mussels are the only freshwater mussel that can attach to objects using sticky threads (byssal threads like those found on the marine blue mussels). These threads allow them to colonize quickly and reach densities of 100,000 or more mussels per square yard. The mussels have an average lifespan of 3.5 to 5 years. A gritty feeling on your boat's hull or other immersed surfaces might indicate that larval zebra mussels have settled.

Zebra mussels originated in the drainage basins of the Black, Caspian, and Aral seas of eastern Europe and have been in western Europe freshwaters since the 1700s. Since first being introduced to North America in 1986, zebra mussels have dramatically altered the balance of freshwater systems and fisheries. These small water dwelling animals have also caused millions of dollars in expenses for industrial water users, drinking water facilities, commercial and recreational boaters, farmers, and other groups and organizations in Canada and the Great Lakes region.

The range occupied by these unwelcome visitors has expanded and continues to grow rapidly. In North America, sightings have been recorded as far north as the Saint Lawrence River near Quebec, as far east as the lower portion of the Hudson River, as far south as the Mississippi River near Vicksburg, and as far west as the Arkansas River in Oklahoma.

In 1993, zebra mussel sightings were confirmed in New England (Lake Champlain). The Lake Champlain population has existed for at least three years, if not longer. Thus, New Hampshire residents and boaters are being encouraged to arm themselves with knowledge about the natural history and geographic spread of the mussels. Interstate boaters and anglers, in particular, should become familiar with boating and fishing practices that decrease the likelihood that zebra mussels will be transferred from an infested water body to an uninfested one.

The infestation risk factor for any particular water body is determined mainly by the amount and type of boat traffic it supports and the chemical characteristics and temperature it maintains. While the goal is to prevent the mussels from becoming established in New England waters, zebra mussels have proven to be adaptable creatures able to survive in a growing range of environmental conditions. Cooperative monitoring activities coordinated by the **New Hampshire Lakes Lay Monitoring Program** will help determine if and when zebra mussels become established in this region. If zebra



mussels are found, information about control techniques can help those concerned choose the best method to reduce the destructive impacts of the mussels.

Take responsibilities for our waters. If you've been boating in fresh water outside of New England within the past 10 days and plan to launch locally, please...

**Inspect** your boat and trailer for weeds. Remove and discard any you find. Zebra mussels are commonly found on aquatic plants in areas of infestation.

**Flush** the cooling system, bilge areas and live wells with tap water.

**Leave** unused bait behind and discard bait bucket water away from surface waters.

**Keep** your boat out of water to dry for 48 hours. If it is visibly fouled by algae, leave it out until the exterior is completely dry or...

**Wash** down the hull at a car wash. Hot (140 degree F) water kills zebra mussels and veligers and high pressure spray helps remove them. Wash fouling off your boat away from water sources!

**Learn** more about the zebra mussel threat in order to be forewarned of the situation and prevent costly repairs or destructive responses.

**Share** information, ideas and monitoring tasks with other members of your lake association, watershed council, marina club, conservation commission, angling group or civic organization.

**Report** any sightings to the **New Hampshire Lakes Lay Monitoring Program**. Preserve specimens in alcohol if possible, note the location where they were found, and send them in to confirm the identification.

To receive more information, request an educational presentation for your next group meeting, become involved in monitoring efforts, or confirm an identification, contact:

**Jeff Schloss**

**Lakes Lay Monitoring Program**

**38 College Road Room 133 Spaulding Hall**

**University of New Hampshire**

**Durham NH 03824-3512**

**(603) 862-3848**

# Understanding Lake Aging (Eutrophication)

by: **Robert Craycraft** Educational Program Coordinator,  
New Hampshire Lakes Lay Monitoring Program  
*University of New Hampshire*  
109 Pettee Hall, Durham, NH 03824  
603-862-3696 FAX: 603-862-0107  
email: bob.craycraft@unh.edu

and **Jeff Schloss** UNH Cooperative Extension Water Resources Specialist

A common concern among **New Hampshire Lakes Lay Monitoring Program (NH LLMP)** participants is a perceived increase in the density and abundance of aquatic plants in the shallows, increases in the amount of microscopic plant “algae” growth (detected as greener water), and water transparency decreases; what is known as **eutrophication**. Eutrophication is a natural process by which all lakes age and progress from clear, pristine lakes to green, nutrient enriched lakes on a geological time frame of thousands of years. Much like the fertilizers applied to our lawns, nutrients that enter our lakes stimulate plant growth and culminate in greener (and in turn less clear) waters. Some lakes age at a faster rate than others due to natural attributes: watershed area relative to lake area, slope of the land surrounding the lake, soil type, mean lake depth, etc. Since our New Hampshire lakes were created during the last ice-age which ended about 10,000 years ago, we should have a natural continuum of lakes ranging from pristine to enriched.

Classification criteria are often used to categorize lakes into what are known as **tro- phic states**, in other words, levels of lake plant and algae productivity or “greenness” Refer to Table 11 below for a summary of commonly used eutrophication parameters.

**Oligotrophic** lakes are considered “unproductive” pristine systems and are characterized by high water clarities, low nutrient concentrations, low algae concentrations, minimal levels of aquatic plant “weed” growth, and high dissolved oxygen concentrations near the lake bottom. **Eutrophic** lakes are considered “highly productive” enriched systems characterized by low water transparencies, high nutrient concentrations,

**Table 11: Eutrophication Parameters and Categorization**

Parameter	Oligotrophic “pristine”	Mesotrophic “transitional”	Eutrophic “enriched”
<b>Chlorophyll a (ug/l) *</b>	<3.0	3.0-7.0	>7.0
<b>Water Transparency (meters) *</b>	>4.0	2.5-4.0	<2.5
<b>Total Phosphorus (ug/l) *</b>	<15.0	15.0-25.0	>25.0
<b>Dissolved Oxygen (saturation) #</b>	high to moderate	moderate to low	low to zero
<b>Macroscopic Plant (Weed) Abundance</b>	low	moderate	high

\* Denotes classification criteria employed by Forsberg and Ryding (1980).

# Denotes dissolved oxygen concentrations near the lakebottom.

high algae concentrations, large stands of aquatic plants and very low dissolved oxygen concentrations near the lake bottom. **Mesotrophic** lakes have qualities between those of oligotrophic and eutrophic lakes and are characterized by moderate water transparencies, moderate nutrient concentrations, moderate algae growth, moderate aquatic plant “weed” growth and decreasing dissolved oxygen concentrations near the lake bottom.

Is a pristine, oligotrophic, lake “better than” an enriched, eutrophic, lake? Not necessarily! As indicated above, lakes will naturally exhibit varying degrees of productivity. Some lakes will naturally be more susceptible to eutrophication than others due to their natural attributes and in turn have aged more rapidly. This is not necessarily a bad thing as our best bass fishing lakes tend to be more mesotrophic to eutrophic than oligotrophic and an ultra-oligotrophic lake (extremely pristine) will not support a very healthy cold water fishery. However, human related activities can augment the aging process (what is known as cultural eutrophication) and result in a transition from a pristine system to an enriched system in tens of years rather than the natural transitional period that should take thousands of years. Cultural eutrophication is particularly a concern for northern New England lakes where large tracts of forested lands are being developed, culminating in an increased susceptibility of these lakes to sediment and nutrient loadings which augment the eutrophication process.

Additionally, other pollutants such as heavy metals, herbicides, insecticides and petroleum products might also affect your lake’s “health”. A “healthy” lake, as far as eutrophication is concerned, is one in which the various aquatic plants and animals are minimally impacted so that nutrients and other materials are processed efficiently. We can liken this process to a well-managed pasture: nutrients stimulate the growth of grasses and other plants that are eaten by grazers like cows and sheep. As long as producers and grazers are balanced, a good amount of nutrients can be processed through the system. Impact the grazers and the grass will overgrow and nuisance weeds will appear, even if nutrients remain the same. In a lake, the producers are the algae and aquatic weeds while the grazers are the microscopic animals (**zooplankton**) and aquatic insects. These organisms can be very susceptible to a wide range of pollutants at very low concentrations. If impacted, the lake can become much more productive and the fishery will be impacted as well since these same organisms are an important food source for most fish at some stage of their life.

Development upon the landscape can negatively affect water quality in a number of ways:

- Removal of shore side vegetation and loss of wetlands - shore side vegetation (what is known as **riparian vegetation**) and wetlands provide a protective buffer that “traps” pollutants before reaching the lake. These buffers remove materials both chemically (through biological uptake) and physically (settling materials out). As riparian buffers are removed and wetlands lost, pollutant materials are more likely to enter the lake and in turn, favor declining water quality.
- Excessive fertilizer applications - fertilizers entering the lake can stimulate aquatic plant and algal growth and in extreme cases result in noxious algal blooms. Increases in algal growth tend to diminish water transparency and under extreme cases culminate in surface “scums” that can wash up on the shoreline and can also produce unpleasant smells as the material decomposes. Exces-

sive nutrient concentrations also favor algal forms known to produce toxins which irritate the skin and under extreme conditions, are dangerous when ingested.

- Increased organic matter loading - organic matter (leaves, grass clippings, etc.) are a major source of nutrients in the aquatic environment. As the vegetative matter decomposes nutrients are “freed up” and can become available for aquatic plant and algal growth. In general, we are not concerned with this material entering the lake naturally (leaf senescence in the fall) but rather excessive loading of this material as occurs when residents dump or rake leaf litter and grass clippings into the lake. This material not only provides large nutrient reserves which can stimulate aquatic plant and algal growth but also makes great habitat for leaches and other potentially undesirable organisms in swimming areas.
- Septic problems - faulty septic systems are a big concern as they can be a primary source of water pollution around our lakes. Septic systems are loaded with nutrients and can also be a health threat when not functioning properly.
- Loss of vegetative cover and the creation of impervious surfaces - A forested watershed offers the best protection against pollutant runoff. Trees and tall vegetation intercept heavy rains that can erode soils and surface materials. The roots of these plants keep the soils in place, process nutrients and absorb moisture so the soils do not wash out. Impervious surfaces (paved roads, parking lots, building roofs, etc.) reduce the water’s capacity to infiltrate into the ground, and in turn, go through nature’s water purification system. As water seeps into the soil, pollutants are removed from the runoff through absorption onto soil particles. Biological processes detoxify substances and/or immobilize substances. Surface water runoff over impervious surfaces also increases water velocities that favor the transport of a greater load of suspended and dissolved pollutants into your lake.

## **How can you minimize your water quality impacts?**

- Minimize fertilizer applications whenever possible. Most people apply far more fertilizers than necessary, with the excess eventually draining into your lake. This not only applies to those immediately adjacent to the lake but to everybody in the watershed. Pollutants in all areas of the watershed will ultimately make their way into your lake. Have your soil tested (the UNH Soils Analytical Laboratory offers soil testing for a nominal fee, contact your county UNH Cooperative Extension Office for further information) to find out how much fertilizer and what type you really need. Sometimes just an application of crushed lime will release enough nutrients to fit the bill. If you do use fertilizer try to use low phosphorus, slow release nitrogen varieties.
- Don’t dump leaf litter or leaves into the lake. Compost the material or take it to a proper waste disposal center. Do not fill in wetland areas. Do not create or enhance beach areas with sand (contains phosphorus, smothers aquatic habitat, fills in lake as it gets transported away by currents and wind).

- Septic systems will not function efficiently without the proper precautionary maintenance. Have your septic system inspected every two to four years and pumped out when necessary. Since the septic system is such an expensive investment often costing around \$10,000 for a complete overhaul, it is advantageous to assure proper care is taken to prolong the system's life. Additionally, following proper maintenance practices will reduce water quality degradation. Refer to:

*Septic Systems, How they work and how to keep them working.* \$1.00/ea  
University of New Hampshire Publications Center (603) 862-2346

*Pipeline: Fall 1995 Vol. 6, No. 4. Maintaining Your Septic System-A Guide for Homeowners.* (\$0.20 ea. plus shipping & handling). 1-800-624-8301

- Maintain shore side (riparian) vegetative cover when new construction is undertaken. For those who have pre-existing houses but lack vegetative buffers, consider shoreline plantings aimed at diminishing the pollution load into your lake. Refer to:

*Planting Shoreland Areas* (no charge) University of New Hampshire Cooperative Extension Publication Center. (603) 862-2346

*A Guide to Developing and Re-Developing Shoreland Property in New Hampshire: A Blueprint to Help You Live by the Water.* North Country Resource Conservation and Development Area, Inc. 103 Main Street-Suite #1, Meredith NH 03253-9266 (603) 279-6546

*Buffers for Wetlands and Surface Waters: A Guidebook for New Hampshire Municipalities.* Audubon Society of New Hampshire. 3 Silk Farm Road, Concord NH 03301 (603) 224-9909 (free for towns, \$5.00 for others).

- If you have shoreland property review the New Hampshire Comprehensive Shoreland Protection Act (CSPA). The CSPA sets legal regulations aimed at protecting water quality. If you have any questions regarding the act or need further information contact the *Shoreline Protection Act Coordinator* at (603) 271-3503.

# Biotoxins: Toxic Chemicals Produced by Cyanobacteria

---

By: James F. Haney and John J. Sasner  
UNH Center for Freshwater Biology

This is an expanded version of an article that appeared in the February 2000 issue of LAKESIDE: a publication of the New Hampshire Lakes Association.

Last summer, George Linscott walked Sam and Magic along the western shore of Lake Champlain. The Labrador retrievers played in the water, but shortly thereafter showed signs of distress. The dogs were rushed to a veterinarian and died later that day. Samples of the water tested indicated the likely cause of the deaths was a toxin produced by *Anabaena*, a type of cyanobacteria, formerly named blue-green algae. High on an alpine meadow in Switzerland cattle die after drinking from what was thought to be a pristine stream. Once again cyanobacteria appeared to be the culprit. These are recent examples of the growing number of cases of cyanobacteria-related deaths of pets and livestock reported worldwide.

Cyanobacteria have been long recognized as problem “algae” in lakes. In addition to imparting bad tastes and odors to water, these microscopic lake inhabitants also produce highly toxic chemicals called biotoxins. Three of the most common toxic cyanobacteria in New Hampshire lakes are *Anabaena*, *Aphanizomenon* and *Microcystis*, commonly referred to as “Annie, Phannie and Mike.” Curiously, their toxicity may change, depending on the lake and time of year and some strains never produce toxins.

Lake biotoxins consist of a variety of chemical compounds. They are often categorized according to their mode of toxicity as (1) neurotoxins, those that interfere with the conduction and transmission of nerve impulses and (2) hepatotoxins, those that cause hemorrhaging of the liver tissue. Whereas the effects of the liver toxins may be seen after hours or days, acute poisoning by neurotoxins may show almost immediate symptoms, such as a tingling sensation followed by paralysis. Both types of biotoxins may be fatal in large doses. Because of widespread damage caused by “red tides”, there is much more known about toxic marine algae. There are, however, surprising similarities between freshwater and marine biotoxins. For example, researchers at the University of New Hampshire discovered that neosaxitoxin, a neurotoxin produced by *Aphanizomenon* (Phannie) is nearly identical to the toxins produced by marine dinoflagellates that cause “red tides” and paralytic shellfish poisons.

Cyanobacteria occur naturally in virtually all lakes. Problems most often occur when excessive nutrients are added to lakes, causing a burst of growth in populations of cyanobacteria and the formation of conspicuous “blooms” along the lakeshore, where they appear as a green film or as blue-green flakes floating just below the surface. When

When toxic, such accumulations pose the greatest health threat if the water containing the bloom is ingested. Direct contact with the skin may cause skin irritations, but does not usually cause serious health effects.

The field of biotoxins research is relatively young and many questions remain unanswered. For example, it is not known what conditions cause certain cyanobacteria to become toxic nor how they may benefit from producing such substances. Since cyanobacteria are among the oldest organisms on earth, these toxins may have evolved long before the presence of other species, such as fish and humans. It is thought that some toxic compounds may in fact be used by cyanobacteria to assist in gathering scarce nutrients necessary for growth, such as iron and carbon.

Using improved methods of detection, a research team from the Biotoxins Lab at the UNH Center for Freshwater Biology, completed its first field season last summer investigating toxins in 50 lakes throughout New Hampshire. The focus of this study was to determine the types of lakes and regions of the state in which the hepatotoxin microcystin are present and also whether these toxins bioaccumulate in the food web of lakes, such as in the crayfish, mussels and fish. Although samples are still being processed and this work has not yet been fully analyzed, preliminary results show that microcystins occur in many lakes throughout the state. Also, the level of microcystins is correlated with the concentration of phosphorus in lake water. This suggests that eutrophication may have a direct linkage to the abundance of biotoxins in our lakes. These findings provide yet another important reason to protect our waters from nutrient pollution and emphasize the need to continually monitor the health of our lakes.

Should New Hampshire lake users be concerned about biotoxins? In most of our lakes concentrations of biotoxins are well below levels that pose a health threat. When swimming it would be wise to avoid areas of the lake where a scum of cyanobacteria have collected. It is especially important to prevent children from playing in such areas, as they may accidentally drink the contaminated lake water. Likewise, pets should be refrained from drinking from areas with conspicuous accumulations of blue-green bacteria.

# REFERENCES

---

- American Public Health Association.(APHA) 1989. Standard Methods for the Examination of Water and Wastewater 17th edition. APHA, AWWA, WPCF.
- Baker, A.L. 1973. Microstratification of phytoplankton in selected Minnesota lakes. Ph. D. thesis, University of Minnesota.
- Carlson, R.E. 1977. A trophic state index for lakes. *Limnol. Oceanogr.* 22:361-379.
- Chase, V.P., L.S. Deming and F. Latawiec. 1995. Buffers for Wetlands and Surface Waters: A Guidebook for New Hampshire Municipalities. Audubon Society of New Hampshire.
- Craycraft, Robert and Jeffrey Schloss. Reports of the NH Lakes Lay Monitoring Program 1990-2001. UNH Center for Freshwater Biology and Cooperative Extension.
- Dates, Geoff and Jeffrey Schloss. 1998. DATA to INFORMATION: Data Management and Analysis for Coastal Monitoring Groups in New Hampshire and Maine-. Maine-NH Sea Grant and Gulf of Maine Council. September 1998. 90 pages. MSG-E-98-8. Manual.
- Edmondson, W.T. 1937. Food conditions in some New Hampshire lakes. In: Biological survey of the Androscoggin, Saco and coastal watersheds. (Report of E.E. Hoover.) New Hampshire Fish and Game Commission. Concord, New Hampshire.
- Estabrook, R.H., J.N. Connor, K.D. Warren, and M.R. Martin. 1987. New Hampshire Lakes and Ponds Inventory. Vol. III. Staff Report No. 153. New Hampshire Department of Environmental Services. Concord, New Hampshire.
- Estabrook, R.H., M.R. Martin and W.M. Henderson. 1988. New Hampshire Lakes and Ponds Inventory. Vol. IV. Staff Report No. 156. New Hampshire Department of Environmental Services. Concord, New Hampshire.
- Estabrook, R.H., M.R. Martin, P.M. McCarthy, D.J. Dubis, and W.M. Henderson. 1989. New Hampshire Lakes and Ponds Inventory. Vol. V. Staff Report No. 166. New Hampshire Department of Environmental Services. Concord, New Hampshire.
- Estabrook, R.H., P.M. McCarthy, M. O'Loan, W.M. Henderson, and D.J. Dubis. 1990. New Hampshire Lakes and Ponds Inventory. Vol. VI. NHDES-WSPCD-90-3. New Hampshire Department of Environmental Services. Concord, New Hampshire.
- Estabrook, R.H., M. O'Loan and W.M. Henderson. 1991. New Hampshire Lakes and Ponds Inventory. Vol. VII. NHDES-WSPCD-91-3. New Hampshire Department of Environmental Services. Concord, New Hampshire.



- Estabrook, R.H., M. O'Loan, W.M. Henderson and K.L. Perkins. 1992. New Hampshire Lakes and Ponds Inventory. Vol. VIII. NHDES-WSPCD-92-6. New Hampshire Department of Environmental Services. Concord, New Hampshire.
- Estabrook, R.H., K. Faul and W.M. Henderson. 1993. New Hampshire Lakes and Ponds Inventory. Vol. IX. NHDES-WSPCD-93-3. New Hampshire Department of Environmental Services. Concord, New Hampshire.
- Estabrook, R.H., K. Faul and W.M. Henderson. 1994. New Hampshire Lakes and Ponds Inventory. Vol. X. NHDES-WSPCD-94-4. New Hampshire Department of Environmental Services. Concord, New Hampshire.
- Estabrook, R.H., W.M. Henderson and S. Ashley. 1996. New Hampshire Lakes and Ponds Inventory. Vol. XII. NHDES-WSPCD-96-6. New Hampshire Department of Environmental Services. Concord, New Hampshire.
- Forsberg, C. and S.O. Ryding. 1980. Eutrophication parameters and trophic state indices in 30 Swedish waste-water receiving lakes. *Arch. Hydrobiol.* 89:189-207
- Gallup, D.N. 1969. Zooplankton distributions and zooplankton-phytoplankton relationships in a mesotrophic lake. Ph.D. Thesis, University of New Hampshire.
- Haney, J.F. and D.J. Hall. 1973. Sugar-coated Daphnia: a preservation technique for Cladocera. *Limnol. Oceanogr.* 18:331-333.
- Hoover, E.E. 1936. Preliminary biological survey of some New Hampshire lakes. Survey report no. 1. New Hampshire Fish and Game Department. Concord, New Hampshire.
- Hoover, E.E. 1937. Biological survey of the Androscoggin, Saco, and coastal watersheds. Survey report no. 2. New Hampshire Fish and Game Department. Concord, New Hampshire.
- Hoover, E.E. 1938. Biological Survey of the Merrimack watershed. Survey report no. 3. New Hampshire Fish and Game Department. Concord, New Hampshire.
- Hutchinson, G.E. 1967. A treatise on limnology, Vol. 2. John Wiley and Sons, New York.
- Lind, O.T. 1979. Handbook of common methods in limnology. C.V. Mosby, St. Louis.
- Lorenzen, M.W. 1980. Use of chlorophyll-Secchi Disk relationships. *Limnol. Oceanogr.* 25:371-372.
- McCafferty, W.P. 1983. Aquatic Entomology: The Fishermen's and Ecologists' Illustrated Guide to Insects and their relatives. Jones and Bartlett Publishers. Boston MA.
- Merritt, R.W. and K.W. Cummins. 1995. An Introduction to the Aquatic Insects of North America. Kendall/Hunt Publishing Company. Dubuque, Iowa

- New Hampshire Water Supply and Pollution Control Commission. 1981. Classification and priority listing of New Hampshire lakes. Vol. II (Parts 1-6). Staff report no. 121. Concord, New Hampshire.
- New Hampshire Water Supply and Pollution Control Commission. 1982. Classification and priority listing of New Hampshire lakes. Vol. III. Staff report no. 121. Concord, New Hampshire.
- New Hampshire Water Supply and Pollution Control Commission. 1983. New Hampshire Lakes and Ponds Inventory. Vol. I. Staff report no. 133. Concord, New Hampshire.
- New Hampshire Water Supply and Pollution Control Commission. 1985. New Hampshire Lakes and Ponds Inventory. Vol. II. Staff report no. 133. Concord, New Hampshire.
- Newell, A.E. 1960. Biological survey of the lakes and ponds in Coos, Grafton and Carroll Counties. Survey report no. 8a. New Hampshire Fish and Game Department. Concord, New Hampshire.
- Newell, A.E. 1970. Biological survey of the lakes and ponds in Cheshire, Hillsborough and Rockingham Counties. Survey report no. 8c. New Hampshire Fish and Game Department. Concord, New Hampshire.
- Newell, A.E. 1977. Biological survey of the lakes and ponds in Sullivan, Merrimack, Belknap and Strafford Counties. Survey report no. 8b. New Hampshire Fish and Game Department. Concord, New Hampshire.
- Schindler, D.W., et al. 1985. Long-term ecosystem stress: Effects of years of experimental acidification on a small lake. *Science*. 228:1395-1400.
- Schloss, Jeffrey A. 2002. GIS Watershed Mapping: Developing and Implementing a Watershed Natural Resources Inventory. In: Handbook of Water Sensitive Ecological Planning & Design- Proceedings of an International Symposium. February 25 - 26, 2000 Harvard University Graduate School of Design. CRC Press, Boca Raton, FL, Chapter II.12.
- Schloss, J. and J. Connor. 2001. Development of Statewide Nutrient (P) Loading Coefficients Through Geographic Information System Aided Analysis. Final Project Report. UNH Water Resources Research Center. Durham, NH.
- Schloss, Jeffrey A. 2000. An Early Success of the Clean Water Action Plan: The Lake Chocorua Project. Proceedings of the 2nd National Water Quality Monitoring Council National Monitoring Conference: Monitoring for the Millennium. April 25-27, 2000 Austin Texas. Environmental Protection Agency, US Geological Survey, Ground Water Protection Council.
- Schloss, Jeffrey A. 1999. Squam Lakes Watershed Natural Resources Inventory: A Case Study for GIS Analysis in a Multijurisdictional Watershed. 1999. Proceedings of the Tijuana River Watershed Workshop. May 5-8, 1999. Tijuana Mexico. NOAA Coastal Services Center publication.


- Schloss, J.A., A.L. Baker and J.F. Haney. 1989. Over a decade of citizen volunteer monitoring in New Hampshire: The New Hampshire Lakes Lay Monitoring Program. Lake and Reservoir Management.
- Sprules, W.G. 1980. Zoogeographic patterns in size structure of zooplankton communities with possible applications to lake ecosystem modeling and management. in W.C. Kerfoot ed. Evolution and Ecology of Zooplankton Communities. University Press of New England. Dartmouth. pp. 642-656.
- Uttermohl, H. 1958. Improvements in the quantitative methods of phytoplankton study. Mitt. int. Ver. Limnol. 9:1-25.
- U.S. Environmental Protection Agency. 1979. A manual of methods for chemical analysis of water and wastes. Office of Technology Transfer, Cincinnati. PA-600/4-79-020.
- Vollenweider, R.A. 1969. A manual on methods for measuring primary productivity in aquatic environments. International Biological Programme. Blackwell Scientific Publications, Oxford.
- Warfel, H.E. 1939. Biological survey of the Connecticut Watershed. Survey Report 4. N.H. Fish and Game. Concord, New Hampshire.
- Wetzel, R.G. 1983. Limnology. Saunders College Publishing, Philadelphia.

# REPORT FIGURES

---


**Figure 11.** Location of the year 2002 Crescent Lake deep sampling station, Sites 6 Center, Wolfeboro New Hampshire.

# Crescent Lake



**Figure 12.** Location of the year 2002 Lake Wentworth deep sampling stations, Sites 1 Fullers, 2 Triggs, and 12 Governors Deep, Wolfeboro New Hampshire.

# Lake Wentworth




**Figure 13.** Crescent Lake, 2002. Seasonal Secchi Disk (water transparency) and chlorophyll *a* trends for Site 6 Center. The Secchi Disk transparency data are reported to the nearest 0.1 meters while the chlorophyll *a* data are reported to the nearest 0.1 parts per billion (ppb).

**Figure 14.** Crescent Lake, 2002. Seasonal Secchi Disk (water transparency) and dissolved color trends for Site 6 Center. The Secchi Disk transparency data are reported to the nearest 0.1 meters while the dissolved color data are reported to the nearest 0.1 chloroplatinate unit (CPU).


Note: the overlay of the Secchi Disk data with chlorophyll *a* and dissolved color data is intended to provide a visual depiction of the impacts of chlorophyll *a* and dissolved color on water transparency measurements (e.g. higher chlorophyll *a* and dissolved color concentrations often correspond to shallower water transparencies).



## Crescent Lake - Site 6 Center (2002 Seasonal Data)



## Crescent Lake - Site 6 Center (2002 Seasonal Data)




**Figure 15.** Lake Wentworth, 2002. Seasonal Secchi Disk (water transparency) and chlorophyll  $a$  trends for Site 1 Fullers. The Secchi Disk transparency data are reported to the nearest 0.1 meters while the chlorophyll  $a$  data are reported to the nearest 0.1 parts per billion (ppb).


**Figure 16.** Lake Wentworth, 2002. Seasonal Secchi Disk (water transparency) and dissolved color trends for Site 1 Fullers. The Secchi Disk transparency data are reported to the nearest 0.1 meters while the dissolved color data are reported to the nearest 0.1 chloroplatinate unit (CPU).

Note: the overlay of the Secchi Disk data with chlorophyll  $a$  and dissolved color data is intended to provide a visual depiction of the impacts of chlorophyll  $a$  and dissolved color on water transparency measurements (e.g. higher chlorophyll  $a$  and dissolved color concentrations often correspond to shallower water transparencies).

## Lake Wentworth - Site 1 Fullers (2002 Seasonal Data)



## Lake Wentworth - Site 1 Fullers (2002 Seasonal Data)




**Figure 17.** Lake Wentworth, 2002. Seasonal Secchi Disk (water transparency) and chlorophyll  $a$  trends for Site 2 Triggs. The Secchi Disk transparency data are reported to the nearest 0.1 meters while the chlorophyll  $a$  data are reported to the nearest 0.1 parts per billion (ppb).


**Figure 18.** Lake Wentworth, 2002. Seasonal Secchi Disk (water transparency) and dissolved color trends for Site 2 Triggs. The Secchi Disk transparency data are reported to the nearest 0.1 meters while the dissolved color data are reported to the nearest 0.1 chloroplatinate unit (CPU).

Note: the overlay of the Secchi Disk data with chlorophyll  $a$  and dissolved color data is intended to provide a visual depiction of the impacts of chlorophyll  $a$  and dissolved color on water transparency measurements (e.g. higher chlorophyll  $a$  and dissolved color concentrations often correspond to shallower water transparencies).

## Lake Wentworth - Site 2 Triggs (2002 Seasonal Data)



## Lake Wentworth - Site 2 Triggs (2002 Seasonal Data)




**Figure 19.** Lake Wentworth, 2002. Seasonal Secchi Disk (water transparency) and chlorophyll  $a$  trends for Site 12 Governors Deep. The Secchi Disk transparency data are reported to the nearest 0.1 meters while the chlorophyll  $a$  data are reported to the nearest 0.1 parts per billion (ppb).


**Figure 20.** Lake Wentworth, 2002. Seasonal Secchi Disk (water transparency) and dissolved color trends for Site 12 Governors Deep. The Secchi Disk transparency data are reported to the nearest 0.1 meters while the dissolved color data are reported to the nearest 0.1 chloroplatinate unit (CPU).

Note: the overlay of the Secchi Disk data with chlorophyll  $a$  and dissolved color data is intended to provide a visual depiction of the impacts of chlorophyll  $a$  and dissolved color on water transparency measurements (e.g. higher chlorophyll  $a$  and dissolved color concentrations often correspond to shallower water transparencies).

## Lake Wentworth - Site 12 Gov. (2002 Seasonal Data)



## Lake Wentworth - Site 12 Gov. (2002 Seasonal Data)




**Figure 21.** Comparison of the year 2002 Crescent Lake, Site 6 Center, lay monitor Secchi Disk transparency data with historical water quality data collected in conjunction with the New Hampshire Lakes Lay Monitoring Program (1984-2001). The shaded regions on the graph denote the ranges characteristic of low, moderate and high Secchi Disk transparencies. The higher the Secchi Disk transparency the clearer the water. Secchi Disk readings are measured to the nearest tenth (0.1) of a meter.


**Figure 22.** Comparison of the year 2002 Crescent Lake, Site 6 Center, lay monitor chlorophyll *a* data with historical water quality data collected in conjunction with the New Hampshire Lakes Lay Monitoring Program (1984-2001). The shaded regions on the graph denote the ranges characteristic of low, moderate and high chlorophyll *a* concentrations. The higher the chlorophyll *a* concentration the greener the water (i.e. more algal growth).



# **CRESCENT LAKE - SITE 6 CENTER** **LAY MONITOR SECCHI DISK TRANSPARENCY DATA** **YEARLY COMPARISONS (1984-2002)**



# **CRESCENT LAKE - SITE 6 CENTER** **LAY MONITOR CHLOROPHYLL *a* DATA** **YEARLY COMPARISONS (1984-2002)**



**Figure 23.** Comparison of the year 2002 Lake Wentworth, Site 1 Fullers, lay monitor Secchi Disk transparency data with historical water quality data collected in conjunction with the New Hampshire Lakes Lay Monitoring Program (1984-2001). The shaded regions on the graph denote the ranges characteristic of low, moderate and high Secchi Disk transparencies. The higher the Secchi Disk transparency the clearer the water. Secchi Disk readings are measured to the nearest tenth (0.1) of a meter.

**Figure 24.** Comparison of the year 2002 Lake Wentworth, Site 1 Fullers, lay monitor chlorophyll  $a$  data with historical water quality data collected in conjunction with the New Hampshire Lakes Lay Monitoring Program (1984-2001). The shaded regions on the graph denote the ranges characteristic of low and moderate chlorophyll  $a$  concentrations. The higher the chlorophyll  $a$  concentration the greener the water (i.e. more algal growth).


**Figure 25.** Comparison of the year 2002 Lake Wentworth, Site 2 Triggs, lay monitor Secchi Disk transparency data with historical water quality data collected in conjunction with the New Hampshire Lakes Lay Monitoring Program (1984-2001). The shaded regions on the graph denote the ranges characteristic of low, moderate and high Secchi Disk transparencies. The higher the Secchi Disk transparency the clearer the water. Secchi Disk readings are measured to the nearest tenth (0.1) of a meter.

**Figure 26.** Comparison of the year 2002 Lake Wentworth, Site 2 Triggs, lay monitor chlorophyll  $\alpha$  data with historical water quality data collected in conjunction with the New Hampshire Lakes Lay Monitoring Program (1984-2001). The shaded regions on the graph denote the ranges characteristic of low and moderate chlorophyll  $\alpha$  concentrations. The higher the chlorophyll  $\alpha$  concentration the greener the water (i.e. more algal growth).

# LAKE WENTWORTH - SITE 2 TRIGGS

## LAY MONITOR SECCHI DISK TRANSPARENCY DATA


### YEARLY COMPARISONS (1985-2002)



# LAKE WENTWORTH - SITE 2 TRIGGS

## LAY MONITOR CHLOROPHYLL a DATA

### YEARLY COMPARISONS (1985-2002)




**Figure 27.** Comparison of the year 2002 Lake Wentworth, Site 12 Governors Deep, lay monitor Secchi Disk transparency data with historical water quality data collected in conjunction with the New Hampshire Lakes Lay Monitoring Program (1987-2001). The shaded regions on the graph denote the ranges characteristic of low, moderate and high Secchi Disk transparencies. The higher the Secchi Disk transparency the clearer the water. Secchi Disk readings are measured to the nearest tenth (0.1) of a meter.

**Figure 28.** Comparison of the year 2002 Lake Wentworth, Site 12 Governors Deep, lay monitor chlorophyll  $a$  data with historical water quality data collected in conjunction with the New Hampshire Lakes Lay Monitoring Program (1987-2001). The shaded regions on the graph denote the ranges characteristic of low and moderate chlorophyll  $a$  concentrations. The higher the chlorophyll  $a$  concentration the greener the water (i.e. more algal growth).

# LAKE WENTWORTH - SITE 12 GOV DEEP

## LAY MONITOR SECCHI DISK TRANSPARENCY DATA


### YEARLY COMPARISONS (1987-2002)



# LAKE WENTWORTH - SITE 12 GOV DEEP

## LAY MONITOR CHLOROPHYLL *a* DATA


### YEARLY COMPARISONS (1987-2002)



**Figure 29.** Temperature and dissolved profiles collected in Crescent Lake, Site 6 Center, on August 1, 2002. The gray shaded region on the graph denotes dissolved oxygen concentrations stressful to warmwater fish. The temperature and dissolved oxygen data were collected at fine increments (generally about 0.2 meters) and are reported as degrees Centigrade ( $^{\circ}\text{C}$ ) and parts per million (ppm), respectively.

# Crescent Lake - Site 6 Center

08/01/02








# APPENDIX A

---

The following graphs depict dissolved oxygen and temperature data collected at the Lake Wentworth deep sampling stations, Sites 1 Fullers, 2 Triggs and 12 Governors Deep, between February 26 and November 1, 2002. Temperature and dissolved oxygen data were collected from the surface down to the lakebottom to discern the degree of temperature stratification and the degree of oxygen depletion near the lakebottom on the respective sampling dates. The temperature units are degrees Centigrade ( $^{\circ}\text{C}$ ) while the dissolved oxygen units are parts per million (ppm). The gray shaded region on the graphs represents dissolved oxygen concentrations stressful to coldwater fish species (dissolved oxygen concentrations less than 5 parts per million). *Notice the decreasing dissolved oxygen concentrations near the lakebottom that occurred between the spring and the fall months.*


# Lake Wentworth - Site 1 Fullers

02/26/02




# Lake Wentworth - Site 1 Fullers

05/09/02




# Lake Wentworth - Site 1 Fullers

05/23/02




# Lake Wentworth - Site 1 Fullers

06/10/02




# Lake Wentworth - Site 1 Fullers

06/29/02




# Lake Wentworth - Site 1 Fullers

07/14/02




# Lake Wentworth - Site 1 Fullers

07/21/02




# Lake Wentworth - Site 1 Fullers

08/01/02




# Lake Wentworth - Site 1 Fullers

## 08/13/02




# Lake Wentworth - Site 1 Fullers

## 08/29/02




# Lake Wentworth - Site 1 Fullers

09/08/02



# Lake Wentworth - Site 1 Fullers


09/22/02






# Lake Wentworth - Site 1 Fullers

## 10/06/02




# Lake Wentworth - Site 1 Fullers

## 11/01/02




# Lake Wentworth - Site 2 Triggs

02/06/02




# Lake Wentworth - Site 2 Triggs

05/09/02




# Lake Wentworth - Site 2 Triggs

05/23/02




# Lake Wentworth - Site 2 Triggs

06/10/02




# Lake Wentworth - Site 2 Triggs

06/29/02




# Lake Wentworth - Site 2 Triggs

07/14/02




# Lake Wentworth - Site 2 Triggs

## 07/21/02




# Lake Wentworth - Site 2 Triggs

## 08/01/02




# Lake Wentworth - Site 2 Triggs

## 08/13/02




# Lake Wentworth - Site 2 Triggs

## 08/29/02




# Lake Wentworth - Site 2 Triggs

09/08/02




# Lake Wentworth - Site 2 Triggs

09/22/02




# Lake Wentworth - Site 2 Triggs

## 10/06/02



# Lake Wentworth - Site 2 Triggs


## 11/01/02






# Lake Wentworth - Site 12 GovDeep

## 02/26/02




# Lake Wentworth - Site 12 GovDeep

## 05/09/02




# Lake Wentworth - Site 12 GovDeep

## 05/23/02




# Lake Wentworth - Site 12 GovDeep

## 06/10/02




# Lake Wentworth - Site 12 GovDeep

## 06/29/02




# Lake Wentworth - Site 12 GovDeep


## 07/14/02



# Lake Wentworth - Site 12 GovDeep 07/21/02




# Lake Wentworth - Site 12 GovDeep 08/01/02




# Lake Wentworth - Site 12 GovDeep

## 08/13/02




# Lake Wentworth - Site 12 GovDeep

## 08/29/02




# Lake Wentworth - Site 12 GovDeep

09/08/02




# Lake Wentworth - Site 12 GovDeep


09/22/02



# Lake Wentworth - Site 12 GovDeep 10/06/02



# Lake Wentworth - Site 12 GovDeep 11/02/02



# APPENDIX B

## Lakes Lay Monitoring Program, U.N.H. [Lay Monitor Data]

Crescent Lake, Wolfeboro New Hampshire  
-- subset of trophic indicators, all sites, 2002

Average transparency:	5.3	(2002:	14 values;	4.7 -	5.9 range)
Average chlorophyll:	1.6	(2002:	14 values;	0.7 -	2.4 range)
Average color:	11.9	(2002:	13 values;	9.6 -	13.9 range)
Average alkalinity (gray):	7.1	(2002:	14 values;	6.9 -	7.9 range)
Average alkalinity (pink):	7.9	(2002:	14 Values;	7.6 -	8.4 range)

Site	Date	Secchi Disk Depth (meters)	Chl <i>a</i> ( $\mu$ g/L)	Dissolved Color (CPU)	Alkalinity gray end pt. @ pH 5.1 (mg/L)	Alkalinity Pink end pt. @ pH 4.6 (mg/L)
6 Center	6/16/02	5.2	2.3	13.0	7.5	8.4
6 Center	6/21/02	5.4	1.5	12.2	6.9	7.9
6 Center	6/28/02	5.9	2.0	13.0	7.9	8.0
6 Center	7/5/02	5.9	0.7	12.2	6.9	7.6
6 Center	7/12/02	5.6	1.3	13.9	7.0	7.6
6 Center	7/19/02	5.1	1.7	10.4	7.1	7.7
6 Center	7/26/02	5.1	1.1	11.3	7.0	7.8
6 Center	8/9/02	5.5	1.5	13.0	6.9	7.6
6 Center	8/16/02	5.0	1.1	9.6	6.9	7.8
6 Center	8/26/02	4.7	2.4	12.2	7.0	8.1
6 Center	8/31/02	5.0	1.9	11.3	7.1	7.9
6 Center	9/6/02	5.0	2.1	11.3	7.1	7.9
6 Center	9/15/02	5.2	2.1	11.3	7.1	7.8
6 Center	9/20/02	5.2	1.3	-----	7.1	7.9

<< End of 2002 data listing, 14 records >>



**Lakes Lay Monitoring Program, U.N.H.**  
**[Lay Monitor Data]**

Wentworth Lake, Wolfeboro New Hampshire  
 -- subset of trophic indicators, all sites, 2002

Average transparency:	7.5	(2002:	31 values;	6.4 - 9.5	range)
Average chlorophyll:	2.0	(2002:	31 values;	0.7 - 5.3	range)
Average color:	11.5	(2002:	33 values;	7.0 - 21.7	range)
Average alkalinity (gray):	6.3	(2002:	33 values;	5.4 - 7.4	range)
Average alkalinity (pink):	7.4	(2002:	33 values;	6.3 - 9.0	range)
Triutary total phos. (ug/L)	10.2	(2002:	27 values;	5.0 - 23.2	range)

Site	Date	Secchi Disk (meters)	Chl <i>a</i> (ppb)	Dissolved Color (CPU)	Alkalinity gray end pt. @ pH 5.1 (mg/L)	Alkalinity pink end pt. @ pH 4.6 (mg/L)	Total Phos. (ug/L)	Specific Conduct. @ 25°C (uS/cm)
1 Fullers	5/8/02	8.0	5.3	13.0	6.4	7.4	-----	-----
1 Fullers	6/25/02	-----	1.7	11.3	7.3	8.0	-----	-----
1 Fullers	6/30/02	8.5	1.3	12.2	6.2	6.7	-----	-----
1 Fullers	7/13/02	7.5	1.8	13.9	6.0	7.0	-----	-----
1 Fullers	7/20/02	7.5	1.7	13.0	6.3	7.3	-----	-----
1 Fullers	7/28/02	7.0	1.9	12.2	6.4	7.1	-----	-----
1 Fullers	8/11/02	9.5	1.3	9.6	7.3	8.3	-----	-----
1 Fullers	8/18/02	7.6	1.7	9.6	6.6	8.4	-----	-----
1 Fullers	8/27/02	-----	-----	9.6	6.4	7.4	-----	-----
1 Fullers	9/1/02	6.5	-----	9.6	6.7	7.1	-----	-----
2 Triggs	6/23/02	7.5	2.2	10.4	5.4	7.9	-----	-----
2 Triggs	6/30/02	7.5	0.8	12.2	7.4	9.0	-----	-----
2 Triggs	7/7/02	7.8	0.7	19.1	5.7	8.2	-----	-----
2 Triggs	7/15/02	7.3	0.9	21.7	6.6	8.8	-----	-----
2 Triggs	7/21/02	7.9	2.4	10.4	6.5	8.7	-----	-----
2 Triggs	7/28/02	7.5	3.0	12.2	6.2	8.4	-----	-----
2 Triggs	8/11/02	7.7	2.3	11.3	6.3	9.0	-----	-----
2 Triggs	8/31/02	6.5	2.6	9.6	5.6	8.0	-----	-----
2 Triggs	9/9/02	8.2	1.9	8.7	5.8	8.0	-----	-----
12 GovDeep	6/14/02	6.6	2.4	13.0	5.8	6.4	-----	-----
12 GovDeep	6/21/02	7.5	2.2	13.0	5.8	6.3	-----	-----
12 GovDeep	6/27/02	7.5	1.3	12.2	5.9	6.5	-----	-----
12 GovDeep	7/7/02	8.7	1.0	13.0	6.2	8.5	-----	-----
12 GovDeep	7/13/02	7.9	1.3	12.2	6.1	6.5	-----	-----
12 GovDeep	7/20/02	7.0	2.4	12.2	6.1	6.4	-----	-----
12 GovDeep	7/28/02	6.6	2.4	9.6	6.0	6.4	-----	-----
12 GovDeep	8/10/02	6.6	2.3	8.7	5.9	6.5	-----	-----
12 GovDeep	8/17/02	8.2	1.8	12.2	5.9	6.5	-----	-----
12 GovDeep	8/25/02	8.1	2.5	8.7	6.3	6.8	-----	-----
12 GovDeep	8/30/02	6.5	2.4	10.4	6.3	6.8	-----	-----
12 GovDeep	9/6/02	6.4	2.6	8.7	6.3	6.8	-----	-----
12 GovDeep	9/13/02	6.9	2.0	7.8	6.3	6.7	-----	-----

Site	Date	Secchi Disk (meters)	Chl <i>a</i> (ppb)	Dissolved Color (CPU)	Alkalinity gray end pt. @ pH 5.1 (mg/L)	Alkalinity pink end pt. @ pH 4.6 (mg/L)	Total Phos. (ug/L)	Specific Conduct. @ 25°C (uS/cm)
12 GovDeep	9/19/02	7.9	2.1	7.0	6.4	6.8	-----	-----
3 Willey Brook	3/27/02	-----	-----	-----	-----	-----	5.3	77.0
3 Willey Brook	11/24/02	-----	-----	-----	-----	-----	7.3	-----
4 Tyler Brook	3/27/02	-----	-----	-----	-----	-----	12.2	74.1
4 Tyler Brook	11/24/02	-----	-----	-----	-----	-----	7.1	-----
5 Ryefield Brk	3/27/02	-----	-----	-----	-----	-----	6.2	51.2
5 Ryefield Brk	11/24/02	-----	-----	-----	-----	-----	8.4	-----
5B Frost Brook	3/27/02	-----	-----	-----	-----	-----	6.5	65.0
5B Frost Brook	11/24/02	-----	-----	-----	-----	-----	12.5	-----
7 Heath Brook	3/27/02	-----	-----	-----	-----	-----	8.3	23.5
7 Heath Brook	11/24/02	-----	-----	-----	-----	-----	6.4	-----
8 Townsend Brk	3/27/02	-----	-----	-----	-----	-----	23.2	62.4
8 Townsend Brk	11/24/02	-----	-----	-----	-----	-----	11.1	-----
9 Warren Brook	3/28/02	-----	-----	-----	-----	-----	7.0	61.9
9 Warren Brook	11/24/02	-----	-----	-----	-----	-----	12.6	-----
11 Fernald Brook	3/27/02	-----	-----	-----	-----	-----	8.5	41.9
11 Fernald Brook	11/24/02	-----	-----	-----	-----	-----	12.0	-----
Breezy Brook	3/27/02	-----	-----	-----	-----	-----	21.5	184.7
Breezy Brook	11/24/02	-----	-----	-----	-----	-----	6.4	-----
Clay Pit Brook (East)	3/27/02	-----	-----	-----	-----	-----	7.5	30.4
Clay Pit Brook (East)	11/24/02	-----	-----	-----	-----	-----	9.7	-----
Clay Pit Brook (West)	3/27/02	-----	-----	-----	-----	-----	15.1	36.5
Clay Pit Brook (West)	11/24/02	-----	-----	-----	-----	-----	19.0	-----
Harvey Brook	3/27/02	-----	-----	-----	-----	-----	8.6	136.0
Harvey Brook	11/24/02	-----	-----	-----	-----	-----	12.5	-----
Outlet	3/27/02	-----	-----	-----	-----	-----	6.6	52.5
Outlet	11/24/02	-----	-----	-----	-----	-----	5.0	-----
Red Brook	3/27/02	-----	-----	-----	-----	-----	9.5	71.6

<< End of 2002 data listing, 60 records >>

**Lakes Lay Monitoring Program, U.N.H.**  
**[Center for Freshwater Biology Data – August 1, 2002]**

Site	Depth (meters)	Chl a (ug/L)	Dissolved Color (CPU)	Carbon Dioxide (mg/L)	Alkalinity gray end point @ pH 5.1 (mg/L)	Alkalinity pink end point @ pH 4.6 (mg/L)	Total Phosphorus (ug/L)
1 Fullers	0.5	1.2	8.7	0.5	6.0	6.5	-----
1 Fullers	3.0	-----	-----	0.6	-----	-----	-----
1 Fullers	10.0	4.5	13.0	3.0	6.2	6.7	8.6
1 Fullers	23.5	-----	-----	5.6	-----	-----	11.2
1 Fullers	0-5.5	1.5	9.6	-----	6.3	6.8	5.7
2 Triggs	0.5	1.2	8.7	0.6	6.0	6.6	-----
2 Triggs	3.0	-----	-----	0.6	-----	-----	-----
2 Triggs	8.0	4.1	8.7	1.7	6.2	6.7	7.8
2 Triggs	14.5	-----	-----	3.4	-----	-----	8.7
2 Triggs	0-7.0	1.9	8.7	-----	6.0	6.6	6.0
6 Center	0.5	1.9	10.4	0.8	6.3	7.1	-----
6 Center	2.0	-----	-----	0.8	-----	-----	-----
6 Center	4.5	-----	-----	0.9	-----	-----	8.4
6 Center	0-3.5	1.3	11.3	-----	6.7	7.2	7.7
12 Governors	0.5	1.9	8.7	0.6	6.1	6.6	-----
12 Governors	3.0	-----	-----	0.6	-----	-----	-----
12 Governors	9.0	4.8	9.6	3.1	6.2	6.8	8.6
12 Governors	18.0	-----	-----	6.3	-----	-----	12.2
12 Governors	0-6.0	1.6	9.6	-----	6.1	6.8	5.8

Site	Secchi Disk Transparency (meters)
1 Fullers	7.6 meters
2 Triggs	7.5 meters
6 Center	5.5 meters (visible on lake bottom)
12 Governors Deep	8.0 meters

Site	Depth (meters)	Temperature (°C)	Dissolved Oxygen (mg/L)	Dissolved Oxygen (% sat.)	Specific Conductivity @ 25°C (uS/cm)	pH (std. units)
1 Fullers	0.08	25.5	7.9	97.1	57.0	6.9
1 Fullers	0.08	25.5	8.0	96.9	57.0	6.9
1 Fullers	0.10	25.6	7.9	97.0	57.0	6.9
1 Fullers	0.24	25.6	7.9	97.1	57.0	6.9
1 Fullers	0.46	25.7	8.0	97.4	57.0	6.9
1 Fullers	0.65	25.7	8.0	97.5	57.0	6.9
1 Fullers	0.83	25.7	7.9	97.3	57.0	6.9
1 Fullers	0.96	25.5	8.0	97.2	57.0	6.9
1 Fullers	1.09	25.3	8.0	97.2	57.0	6.9
1 Fullers	1.27	25.1	8.0	97.4	57.0	6.9
1 Fullers	1.41	25.0	8.1	97.4	57.0	6.9
1 Fullers	1.57	24.9	8.1	97.5	57.0	6.9

Site	Depth (meters)	Temperature (°C)	Dissolved Oxygen (mg/L)	Dissolved Oxygen (% sat.)	Specific Conductivity @ 25°C (uS/cm)	pH (std. units)
1 Fullers	1.75	24.9	8.1	97.5	57.0	6.9
1 Fullers	1.92	24.8	8.1	97.4	57.0	6.9
1 Fullers	2.09	24.8	8.1	97.4	57.0	6.9
1 Fullers	2.24	24.8	8.1	97.4	57.0	6.9
1 Fullers	2.42	24.8	8.1	97.3	57.0	6.9
1 Fullers	2.61	24.8	8.1	97.3	57.0	6.9
1 Fullers	2.86	24.8	8.1	97.3	57.0	6.9
1 Fullers	3.06	24.7	8.1	97.1	57.0	6.9
1 Fullers	3.20	24.7	8.1	97.1	57.0	6.9
1 Fullers	3.38	24.7	8.1	97.1	57.0	6.9
1 Fullers	3.57	24.7	8.1	97.2	57.0	6.9
1 Fullers	3.78	24.7	8.1	97.2	57.0	6.9
1 Fullers	3.95	24.7	8.1	97.0	57.0	6.9
1 Fullers	4.11	24.7	8.1	97.0	57.0	6.9
1 Fullers	4.26	24.6	8.1	97.0	57.0	6.9
1 Fullers	4.47	24.6	8.1	97.0	57.0	6.9
1 Fullers	4.72	24.6	8.1	96.6	57.0	6.9
1 Fullers	5.08	24.5	8.0	95.7	57.0	6.9
1 Fullers	5.43	24.3	7.9	94.7	57.0	6.9
1 Fullers	5.66	24.1	7.9	93.4	57.0	6.9
1 Fullers	5.86	23.8	7.8	92.0	57.0	6.9
1 Fullers	6.03	23.6	7.7	90.8	57.0	6.9
1 Fullers	6.19	23.4	7.7	90.1	57.0	6.9
1 Fullers	6.35	23.3	7.7	89.6	57.0	6.9
1 Fullers	6.50	23.2	7.6	89.4	57.0	6.9
1 Fullers	6.66	23.1	7.6	88.8	57.0	6.8
1 Fullers	6.81	22.9	7.6	88.1	57.0	6.8
1 Fullers	6.94	22.6	7.5	86.8	57.0	6.8
1 Fullers	7.14	22.4	7.4	84.9	57.0	6.8
1 Fullers	7.38	22.2	7.3	83.2	57.0	6.8
1 Fullers	7.52	21.9	7.1	80.8	57.0	6.8
1 Fullers	7.65	21.7	6.9	78.7	57.0	6.8
1 Fullers	7.82	21.5	6.8	77.5	57.0	6.8
1 Fullers	7.98	21.4	6.8	76.4	57.0	6.7
1 Fullers	8.17	21.3	6.7	75.6	57.0	6.7
1 Fullers	8.33	21.2	6.7	74.8	57.0	6.7
1 Fullers	8.49	21.0	6.6	73.8	57.0	6.7
1 Fullers	8.64	20.9	6.5	72.5	57.0	6.7
1 Fullers	8.79	20.6	6.4	70.9	57.0	6.7
1 Fullers	8.91	20.3	6.2	68.9	57.0	6.6
1 Fullers	9.10	20.0	6.1	66.6	57.0	6.6
1 Fullers	9.24	19.8	5.9	64.4	57.0	6.6
1 Fullers	9.33	19.6	5.7	62.2	57.0	6.6
1 Fullers	9.45	19.4	5.5	60.2	56.0	6.6
1 Fullers	9.60	19.2	5.4	58.6	56.0	6.6
1 Fullers	9.78	19.0	5.3	57.3	56.0	6.5

Site	Depth (meters)	Temperature (°C)	Dissolved Oxygen (mg/L)	Dissolved Oxygen (% sat.)	Specific Conductivity @ 25°C (uS/cm)	pH (std. units)
1 Fullers	9.95	18.8	5.2	55.5	56.0	6.5
1 Fullers	10.08	18.5	5.0	53.3	56.0	6.5
1 Fullers	10.20	18.3	4.8	51.2	56.0	6.5
1 Fullers	10.33	18.1	4.7	49.9	56.0	6.5
1 Fullers	10.47	18.0	4.6	48.7	56.0	6.5
1 Fullers	10.61	17.9	4.5	47.5	56.0	6.5
1 Fullers	10.79	17.8	4.4	46.7	56.0	6.4
1 Fullers	10.98	17.7	4.4	45.7	56.0	6.4
1 Fullers	11.15	17.6	4.3	44.6	56.0	6.4
1 Fullers	11.31	17.4	4.2	43.4	56.0	6.4
1 Fullers	11.50	17.2	4.1	42.1	56.0	6.4
1 Fullers	11.64	17.0	4.0	40.8	56.0	6.4
1 Fullers	11.80	16.6	3.8	39.1	56.0	6.4
1 Fullers	11.96	16.3	3.7	38.0	56.0	6.4
1 Fullers	12.13	16.0	3.6	36.7	56.0	6.3
1 Fullers	12.28	15.7	3.5	35.5	56.0	6.3
1 Fullers	12.47	15.5	3.5	34.8	56.0	6.3
1 Fullers	12.64	15.3	3.5	34.8	56.0	6.3
1 Fullers	12.85	15.1	3.5	34.4	56.0	6.3
1 Fullers	13.10	14.9	3.4	33.8	55.0	6.3
1 Fullers	13.28	14.6	3.4	33.7	55.0	6.3
1 Fullers	13.46	14.4	3.4	33.5	55.0	6.3
1 Fullers	13.62	14.2	3.4	33.5	55.0	6.3
1 Fullers	13.83	14.1	3.4	33.2	55.0	6.2
1 Fullers	14.01	14.0	3.4	32.7	55.0	6.2
1 Fullers	14.22	13.8	3.4	32.4	55.0	6.2
1 Fullers	14.45	13.6	3.4	32.4	55.0	6.2
1 Fullers	14.64	13.5	3.3	32.0	55.0	6.2
1 Fullers	14.88	13.3	3.3	31.2	55.0	6.2
1 Fullers	15.06	13.1	3.3	31.2	55.0	6.2
1 Fullers	15.25	12.9	3.3	31.1	55.0	6.2
1 Fullers	15.45	12.8	3.3	31.0	55.0	6.2
1 Fullers	15.63	12.7	3.3	30.8	55.0	6.2
1 Fullers	15.79	12.7	3.2	30.4	55.0	6.1
1 Fullers	15.92	12.6	3.2	30.1	55.0	6.1
1 Fullers	16.12	12.6	3.2	30.0	55.0	6.1
1 Fullers	16.36	12.6	3.2	30.0	55.0	6.1
1 Fullers	16.56	12.6	3.2	29.7	54.0	6.1
1 Fullers	16.74	12.5	3.1	29.4	54.0	6.1
1 Fullers	16.95	12.5	3.1	29.1	54.0	6.1
1 Fullers	17.13	12.5	3.1	28.6	54.0	6.1
1 Fullers	17.33	12.5	3.0	28.4	54.0	6.1
1 Fullers	17.51	12.5	3.0	28.1	54.0	6.1
1 Fullers	17.69	12.4	3.0	27.9	54.0	6.1
1 Fullers	17.87	12.4	3.0	27.7	54.0	6.1
1 Fullers	18.05	12.4	2.9	27.3	54.0	6.1

Site	Depth (meters)	Temperature (°C)	Dissolved Oxygen (mg/L)	Dissolved Oxygen (% sat.)	Specific Conductivity @ 25°C (uS/cm)	pH (std. units)
1 Fullers	18.23	12.4	2.9	26.9	54.0	6.1
1 Fullers	18.40	12.4	2.9	26.7	54.0	6.1
1 Fullers	18.59	12.4	2.8	26.5	54.0	6.0
1 Fullers	18.75	12.3	2.8	26.2	54.0	6.0
1 Fullers	18.92	12.3	2.8	25.9	54.0	6.0
1 Fullers	19.08	12.3	2.8	25.7	54.0	6.0
1 Fullers	19.30	12.3	2.7	25.4	54.0	6.0
1 Fullers	19.48	12.3	2.7	25.1	54.0	6.0
1 Fullers	19.65	12.3	2.7	24.8	54.0	6.0
1 Fullers	19.85	12.3	2.6	24.4	54.0	6.0
1 Fullers	20.03	12.2	2.6	24.2	54.0	6.0
1 Fullers	20.22	12.2	2.6	24.0	54.0	6.0
1 Fullers	20.38	12.2	2.6	23.9	54.0	6.0
1 Fullers	20.55	12.2	2.6	23.8	54.0	6.0
1 Fullers	20.85	12.2	2.6	23.8	54.0	6.0
1 Fullers	21.15	12.2	2.5	23.7	54.0	6.0
1 Fullers	21.38	12.2	2.5	23.6	54.0	6.0
1 Fullers	21.64	12.2	2.5	23.5	54.0	6.0
1 Fullers	21.84	12.2	2.5	23.5	54.0	6.0
1 Fullers	21.99	12.2	2.5	23.4	54.0	6.0
1 Fullers	22.17	12.2	2.5	23.3	54.0	6.0
1 Fullers	22.33	12.2	2.5	23.1	55.0	6.0
1 Fullers	22.48	12.2	2.5	22.9	55.0	6.0
1 Fullers	22.69	12.2	2.4	22.7	55.0	6.0
1 Fullers	22.91	12.2	2.4	22.6	55.0	5.9
1 Fullers	23.14	12.2	2.4	22.4	55.0	5.9
1 Fullers	23.36	12.2	2.4	22.0	55.0	5.9
1 Fullers	23.61	12.2	2.3	21.6	55.0	5.9
1 Fullers	23.90	12.1	2.3	21.2	55.0	5.9
1 Fullers	24.16	12.1	2.3	20.9	55.0	5.9
1 Fullers	24.36	12.1	2.2	20.3	55.0	5.9
2 Triggs	0.09	25.9	8.0	98.5	57.0	7.0
2 Triggs	0.11	26.1	8.0	98.9	56.0	7.0
2 Triggs	0.24	26.0	8.1	99.2	56.0	7.0
2 Triggs	0.37	25.8	8.1	99.1	56.0	7.0
2 Triggs	0.60	25.5	8.1	99.1	56.0	7.0
2 Triggs	0.88	25.3	8.2	99.8	57.0	7.0
2 Triggs	1.13	25.1	8.2	99.6	57.0	7.0
2 Triggs	1.39	25.0	8.2	99.4	57.0	7.0
2 Triggs	1.60	24.9	8.3	100.1	57.0	7.0
2 Triggs	1.77	24.9	8.3	100.5	57.0	7.0
2 Triggs	1.93	24.9	8.3	100.0	57.0	7.0
2 Triggs	2.12	24.9	8.2	99.2	57.0	7.0
2 Triggs	2.27	24.9	8.3	99.7	57.0	7.0
2 Triggs	2.40	24.8	8.3	99.7	57.0	7.0
2 Triggs	2.58	24.8	8.2	99.2	57.0	7.0

Site	Depth (meters)	Temperature (°C)	Dissolved Oxygen (mg/L)	Dissolved Oxygen (% sat.)	Specific Conductivity @ 25°C (uS/cm)	pH (std. units)
2 Triggs	2.74	24.8	8.2	98.7	57.0	7.0
2 Triggs	2.90	24.8	8.2	98.8	57.0	7.0
2 Triggs	3.04	24.8	8.2	98.8	57.0	7.0
2 Triggs	3.16	24.8	8.2	98.5	57.0	7.0
2 Triggs	3.34	24.8	8.2	98.3	57.0	7.0
2 Triggs	3.50	24.8	8.2	98.4	57.0	7.0
2 Triggs	3.68	24.8	8.2	98.5	57.0	7.0
2 Triggs	3.81	24.8	8.2	98.5	57.0	7.0
2 Triggs	3.93	24.7	8.2	98.4	57.0	7.0
2 Triggs	4.05	24.7	8.2	98.4	57.0	7.0
2 Triggs	4.16	24.7	8.2	98.2	56.0	7.0
2 Triggs	4.29	24.7	8.1	97.9	57.0	7.0
2 Triggs	4.44	24.7	8.1	97.8	57.0	7.0
2 Triggs	4.57	24.7	8.1	97.7	57.0	7.0
2 Triggs	4.71	24.7	8.1	97.6	57.0	7.0
2 Triggs	4.89	24.7	8.1	97.4	57.0	7.0
2 Triggs	5.02	24.6	8.1	97.3	57.0	7.0
2 Triggs	5.19	24.6	8.1	96.9	57.0	7.0
2 Triggs	5.34	24.6	8.1	96.9	56.0	7.0
2 Triggs	5.47	24.6	8.1	96.7	56.0	7.0
2 Triggs	5.62	24.6	8.1	96.7	56.0	7.0
2 Triggs	5.76	24.6	8.1	96.7	56.0	7.0
2 Triggs	5.91	24.5	8.0	96.3	56.0	7.0
2 Triggs	6.07	24.4	8.0	95.8	56.0	7.0
2 Triggs	6.22	24.3	8.0	95.4	57.0	6.9
2 Triggs	6.39	24.2	7.9	94.2	57.0	6.9
2 Triggs	6.54	24.1	7.9	93.6	57.0	6.9
2 Triggs	6.69	24.0	7.9	93.2	57.0	6.9
2 Triggs	6.80	23.9	7.8	92.8	57.0	6.9
2 Triggs	6.91	23.9	7.8	92.1	57.0	6.9
2 Triggs	7.01	23.8	7.7	91.1	57.0	6.9
2 Triggs	7.11	23.6	7.6	89.8	57.0	6.9
2 Triggs	7.22	23.5	7.5	88.7	57.0	6.9
2 Triggs	7.35	23.3	7.5	87.6	57.0	6.9
2 Triggs	7.51	23.1	7.4	86.3	57.0	6.9
2 Triggs	7.65	22.9	7.3	85.1	57.0	6.9
2 Triggs	7.82	22.8	7.3	84.5	57.0	6.8
2 Triggs	7.98	22.6	7.2	83.2	57.0	6.8
2 Triggs	8.16	22.4	7.1	81.6	57.0	6.8
2 Triggs	8.34	22.2	7.0	79.8	57.0	6.8
2 Triggs	8.48	22.1	6.8	78.2	57.0	6.8
2 Triggs	8.63	21.9	6.7	76.5	57.0	6.8
2 Triggs	8.77	21.7	6.6	74.5	57.0	6.8
2 Triggs	8.94	21.6	6.4	72.2	57.0	6.7
2 Triggs	9.09	21.4	6.2	69.9	57.0	6.7
2 Triggs	9.26	21.3	6.0	67.9	57.0	6.7

Site	Depth (meters)	Temperature (°C)	Dissolved Oxygen (mg/L)	Dissolved Oxygen (% sat.)	Specific Conductivity @ 25°C (uS/cm)	pH (std. units)
2 Triggs	9.44	21.2	5.9	66.5	57.0	6.7
2 Triggs	9.59	21.2	5.8	65.4	57.0	6.7
2 Triggs	9.74	21.1	5.7	64.4	57.0	6.7
2 Triggs	9.90	21.0	5.6	63.2	57.0	6.6
2 Triggs	10.05	21.0	5.6	62.2	57.0	6.6
2 Triggs	10.21	20.9	5.5	61.5	57.0	6.6
2 Triggs	10.38	20.9	5.5	61.0	57.0	6.6
2 Triggs	10.58	20.9	5.4	60.6	57.0	6.6
2 Triggs	10.78	20.9	5.4	60.4	57.0	6.6
2 Triggs	10.99	20.9	5.4	60.1	56.0	6.6
2 Triggs	11.18	20.9	5.4	60.0	56.0	6.6
2 Triggs	11.35	20.9	5.3	59.7	56.0	6.6
2 Triggs	11.51	20.9	5.3	59.4	56.0	6.5
2 Triggs	11.69	20.9	5.3	59.2	56.0	6.5
2 Triggs	11.85	20.8	5.3	59.0	56.0	6.5
2 Triggs	12.03	20.8	5.3	58.7	56.0	6.5
2 Triggs	12.22	20.8	5.2	58.6	56.0	6.5
2 Triggs	12.43	20.8	5.2	58.3	56.0	6.5
2 Triggs	12.63	20.8	5.2	57.7	56.0	6.5
2 Triggs	12.79	20.8	5.1	57.1	56.0	6.5
2 Triggs	12.93	20.7	5.1	56.7	56.0	6.5
2 Triggs	13.08	20.7	5.1	56.6	56.0	6.5
2 Triggs	13.28	20.7	5.1	56.5	56.0	6.5
2 Triggs	13.46	20.7	5.0	56.2	56.0	6.5
2 Triggs	13.60	20.6	5.0	55.4	56.0	6.4
2 Triggs	13.71	20.6	4.9	54.8	56.0	6.4
2 Triggs	13.81	20.5	4.9	54.6	57.0	6.4
2 Triggs	13.95	20.5	4.9	54.2	57.0	6.4
2 Triggs	14.12	20.3	4.8	52.6	57.0	6.4
2 Triggs	14.27	20.1	4.5	50.0	57.0	6.4
2 Triggs	14.41	20.0	4.2	46.4	57.0	6.4
2 Triggs	14.59	19.9	3.9	43.1	57.0	6.4
2 Triggs	14.76	19.7	3.6	39.8	58.0	6.4
2 Triggs	14.92	19.4	3.3	35.6	59.0	6.4
2 Triggs	15.09	19.1	2.7	29.4	59.0	6.4
2 Triggs	15.23	18.9	2.1	22.4	60.0	6.3
2 Triggs	15.33	18.7	1.7	17.6	61.0	6.3
2 Triggs	15.41	18.6	1.3	13.9	61.0	6.3
6 Center	0.06	27.1	7.9	99.2	59.0	6.8
6 Center	0.07	27.2	7.9	99.2	59.0	6.8
6 Center	0.23	27.3	7.9	99.4	59.0	6.8
6 Center	0.44	27.2	7.9	99.4	59.0	6.8
6 Center	0.75	27.1	7.9	99.7	59.0	6.8
6 Center	1.09	26.8	8.0	99.7	59.0	6.8
6 Center	1.50	26.6	8.0	100.0	59.0	6.8
6 Center	1.90	26.5	8.0	99.8	59.0	6.8



Site	Depth (meters)	Temperature (°C)	Dissolved Oxygen (mg/L)	Dissolved Oxygen (% sat.)	Specific Conductivity @ 25°C (uS/cm)	pH (std. units)
6 Center	2.26	26.4	8.0	99.7	59.0	6.8
6 Center	2.67	26.3	8.0	99.5	59.0	6.8
6 Center	3.18	26.1	8.0	98.6	59.0	6.8
6 Center	3.60	25.7	7.9	97.4	59.0	6.8
6 Center	3.97	25.3	8.1	98.1	59.0	6.8
6 Center	4.27	24.9	8.2	99.3	59.0	6.8
6 Center	4.53	24.5	8.3	99.4	59.0	6.8
6 Center	4.83	24.2	8.3	98.9	60.0	6.8
6 Center	5.13	23.7	8.2	97.4	62.0	6.8
6 Center	5.31	23.2	7.7	89.7	63.0	6.7
12 Governors	0.25	25.5	8.0	97.8	56.0	7.2
12 Governors	0.26	25.6	8.0	98.2	56.0	7.2
12 Governors	0.34	25.7	8.0	98.2	56.0	7.2
12 Governors	0.42	25.7	8.0	98.0	56.0	7.2
12 Governors	0.42	25.5	8.0	98.0	56.0	7.2
12 Governors	0.51	25.6	8.0	98.0	56.0	7.2
12 Governors	0.59	25.5	8.1	98.3	56.0	7.2
12 Governors	0.62	25.6	8.0	97.9	56.0	7.2
12 Governors	0.75	25.5	8.0	97.8	56.0	7.2
12 Governors	0.90	25.4	8.0	97.8	56.0	7.2
12 Governors	1.00	25.4	8.0	97.9	56.0	7.2
12 Governors	1.11	25.4	8.0	97.7	56.0	7.2
12 Governors	1.17	25.3	8.0	97.6	56.0	7.2
12 Governors	1.22	25.3	8.0	97.7	56.0	7.2
12 Governors	1.33	25.3	8.0	97.6	56.0	7.2
12 Governors	1.46	25.3	8.0	97.7	56.0	7.2
12 Governors	1.62	25.3	8.0	97.6	56.0	7.2
12 Governors	1.75	25.3	8.0	97.7	56.0	7.2
12 Governors	1.90	25.2	8.0	97.6	56.0	7.2
12 Governors	2.08	25.2	8.0	97.5	56.0	7.2
12 Governors	2.24	25.2	8.0	97.5	56.0	7.2
12 Governors	2.36	25.2	8.0	97.3	56.0	7.2
12 Governors	2.48	25.2	8.0	97.2	56.0	7.2
12 Governors	2.63	25.1	8.0	97.1	56.0	7.2
12 Governors	2.77	25.1	8.0	97.1	56.0	7.2
12 Governors	2.92	25.1	8.0	97.1	56.0	7.2
12 Governors	3.10	25.1	8.0	97.1	56.0	7.2
12 Governors	3.27	25.1	8.0	97.2	56.0	7.2
12 Governors	3.49	25.1	8.0	97.2	56.0	7.2
12 Governors	3.70	25.1	8.0	97.3	56.0	7.2
12 Governors	3.87	25.1	8.0	97.1	56.0	7.2
12 Governors	4.07	25.1	8.0	97.0	56.0	7.2
12 Governors	4.23	25.1	8.0	97.0	56.0	7.2
12 Governors	4.37	25.1	8.0	97.0	56.0	7.2
12 Governors	4.54	25.1	8.0	97.1	56.0	7.2
12 Governors	4.67	25.1	8.0	97.0	56.0	7.2

Site	Depth (meters)	Temperature (°C)	Dissolved Oxygen (mg/L)	Dissolved Oxygen (% sat.)	Specific Conductivity @ 25°C (uS/cm)	pH (std. units)
12 Governors	4.81	25.1	8.0	96.9	56.0	7.2
12 Governors	4.93	25.0	8.0	96.9	56.0	7.2
12 Governors	5.08	25.0	8.0	96.9	56.0	7.2
12 Governors	5.21	25.0	8.0	97.0	56.0	7.2
12 Governors	5.36	25.0	8.0	96.9	56.0	7.2
12 Governors	5.53	25.0	8.0	96.7	56.0	7.2
12 Governors	5.69	25.0	8.0	96.6	56.0	7.2
12 Governors	5.81	25.0	8.0	96.5	56.0	7.2
12 Governors	5.97	24.9	8.0	96.4	56.0	7.2
12 Governors	6.08	24.8	8.0	96.4	56.0	7.2
12 Governors	6.19	24.7	8.0	96.1	56.0	7.2
12 Governors	6.26	24.6	8.0	96.0	56.0	7.2
12 Governors	6.37	24.4	8.0	95.4	56.0	7.2
12 Governors	6.48	24.3	7.9	94.8	57.0	7.2
12 Governors	6.59	24.1	7.9	94.2	57.0	7.2
12 Governors	6.69	24.0	7.8	93.1	57.0	7.2
12 Governors	6.80	23.9	7.8	92.3	56.0	7.2
12 Governors	6.89	23.7	7.7	90.5	56.0	7.1
12 Governors	7.02	23.5	7.5	88.7	56.0	7.1
12 Governors	7.15	23.4	7.4	87.2	56.0	7.1
12 Governors	7.24	23.2	7.3	85.2	56.0	7.1
12 Governors	7.33	23.0	7.1	83.1	56.0	7.1
12 Governors	7.44	22.9	7.0	81.7	57.0	7.1
12 Governors	7.53	22.7	6.9	80.3	57.0	7.1
12 Governors	7.63	22.5	6.9	79.0	57.0	7.1
12 Governors	7.74	22.3	6.8	78.0	57.0	7.0
12 Governors	7.87	22.0	6.7	76.4	57.0	7.0
12 Governors	8.02	21.7	6.5	74.2	57.0	7.0
12 Governors	8.15	21.5	6.4	72.1	57.0	7.0
12 Governors	8.29	21.3	6.3	70.6	57.0	7.0
12 Governors	8.43	21.1	6.2	69.4	57.0	6.9
12 Governors	8.56	21.0	6.0	67.6	57.0	6.9
12 Governors	8.72	20.9	5.9	66.4	56.0	6.9
12 Governors	8.85	20.7	5.9	65.3	57.0	6.9
12 Governors	8.96	20.5	5.8	64.2	56.0	6.9
12 Governors	9.07	20.4	5.7	63.4	56.0	6.9
12 Governors	9.18	20.3	5.7	62.6	56.0	6.9
12 Governors	9.31	20.1	5.6	61.6	56.0	6.8
12 Governors	9.44	20.0	5.5	60.4	56.0	6.8
12 Governors	9.61	19.8	5.4	59.0	56.0	6.8
12 Governors	9.77	19.7	5.3	58.1	56.0	6.8
12 Governors	9.92	19.6	5.3	57.4	56.0	6.8
12 Governors	10.12	19.3	5.2	56.6	56.0	6.8
12 Governors	10.28	18.9	5.1	55.2	56.0	6.8
12 Governors	10.43	18.5	5.0	53.0	56.0	6.7
12 Governors	10.55	18.2	4.7	50.0	56.0	6.7

Site	Depth (meters)	Temperature (°C)	Dissolved Oxygen (mg/L)	Dissolved Oxygen (% sat.)	Specific Conductivity @ 25°C (uS/cm)	pH (std. units)
12 Governors	10.68	17.9	4.5	47.3	56.0	6.7
12 Governors	10.83	17.6	4.3	45.4	56.0	6.7
12 Governors	10.97	17.4	4.2	43.7	56.0	6.7
12 Governors	11.11	17.2	4.1	42.4	56.0	6.7
12 Governors	11.21	17.0	4.0	41.1	56.0	6.7
12 Governors	11.31	16.7	3.9	39.8	56.0	6.7
12 Governors	11.43	16.4	3.8	38.7	56.0	6.6
12 Governors	11.56	16.2	3.7	37.7	56.0	6.6
12 Governors	11.67	16.0	3.6	36.8	56.0	6.6
12 Governors	11.78	15.9	3.6	36.2	56.0	6.6
12 Governors	11.89	15.8	3.5	35.5	56.0	6.6
12 Governors	12.01	15.7	3.5	35.4	56.0	6.6
12 Governors	12.16	15.5	3.5	34.6	56.0	6.6
12 Governors	12.33	15.2	3.4	34.1	56.0	6.6
12 Governors	12.47	15.0	3.4	33.3	56.0	6.5
12 Governors	12.60	14.8	3.3	32.9	56.0	6.5
12 Governors	12.76	14.6	3.3	32.5	56.0	6.5
12 Governors	12.86	14.5	3.3	32.1	56.0	6.5
12 Governors	12.98	14.3	3.2	31.6	56.0	6.5
12 Governors	13.12	14.1	3.2	31.3	56.0	6.5
12 Governors	13.22	14.0	3.2	30.8	55.0	6.5
12 Governors	13.33	13.8	3.2	30.5	55.0	6.5
12 Governors	13.46	13.7	3.1	30.3	55.0	6.5
12 Governors	13.60	13.5	3.1	29.9	55.0	6.4
12 Governors	13.80	13.4	3.1	29.2	55.0	6.4
12 Governors	14.04	13.3	3.0	28.7	55.0	6.4
12 Governors	14.24	13.2	2.9	27.6	55.0	6.4
12 Governors	14.43	13.2	2.8	26.9	55.0	6.4
12 Governors	14.59	13.2	2.8	26.4	55.0	6.4
12 Governors	14.73	13.1	2.7	25.9	55.0	6.4
12 Governors	14.89	13.0	2.7	25.2	55.0	6.4
12 Governors	15.05	12.9	2.6	24.4	56.0	6.4
12 Governors	15.23	12.8	2.5	23.5	56.0	6.3
12 Governors	15.43	12.7	2.4	22.5	56.0	6.3
12 Governors	15.65	12.7	2.3	21.4	56.0	6.3
12 Governors	15.86	12.6	2.2	20.6	56.0	6.3
12 Governors	16.03	12.6	2.1	20.0	56.0	6.3
12 Governors	16.18	12.6	2.1	19.6	56.0	6.3
12 Governors	16.36	12.6	2.1	19.4	56.0	6.3
12 Governors	16.54	12.6	2.0	19.2	56.0	6.3
12 Governors	16.74	12.6	2.0	18.5	56.0	6.3
12 Governors	16.90	12.5	1.9	17.8	56.0	6.3
12 Governors	17.09	12.5	1.8	17.1	56.0	6.3
12 Governors	17.24	12.5	1.8	16.6	56.0	6.3
12 Governors	17.43	12.5	1.7	16.2	56.0	6.3
12 Governors	17.61	12.5	1.7	15.8	56.0	6.3

Site	Depth (meters)	Temperature (°C)	Dissolved Oxygen (mg/L)	Dissolved Oxygen (% sat.)	Specific Conductivity @ 25°C (uS/cm)	pH (std. units)
12 Governors	17.81	12.5	1.7	15.6	56.0	6.2
12 Governors	18.00	12.5	1.6	15.4	56.0	6.2
12 Governors	18.19	12.5	1.6	15.3	56.0	6.2
12 Governors	18.45	12.5	1.6	15.2	56.0	6.2
12 Governors	18.77	12.5	1.6	15.2	56.0	6.2
12 Governors	19.00	12.5	1.6	15.1	56.0	6.2
12 Governors	19.25	12.5	1.6	14.9	56.0	6.2
12 Governors	19.53	12.5	1.6	14.8	56.0	6.2
12 Governors	19.82	12.5	1.6	14.6	56.0	6.2
12 Governors	20.00	12.5	1.5	14.5	56.0	6.2
12 Governors	20.14	12.5	1.5	14.4	56.0	6.2
12 Governors	20.29	12.5	1.5	14.3	56.0	6.2
12 Governors	20.43	12.5	1.5	14.3	56.0	6.2
12 Governors	20.59	12.5	1.5	14.2	56.0	6.2
12 Governors	20.73	12.5	1.5	14.1	56.0	6.2
12 Governors	20.87	12.5	1.5	14.1	56.0	6.2
12 Governors	20.95	12.5	1.5	14.0	56.0	6.2
12 Governors	20.99	12.5	1.5	13.8	56.0	6.2
12 Governors	21.04	12.5	1.5	13.7	56.0	6.2
12 Governors	21.07	12.5	1.5	13.6	56.0	6.2
12 Governors	21.10	12.5	1.4	13.4	56.0	6.2
12 Governors	21.11	12.5	1.4	13.4	56.0	6.1
12 Governors	21.12	12.5	1.4	13.3	56.0	6.1
12 Governors	21.12	12.5	1.4	13.2	56.0	6.1



# APPENDIX C

---

## GLOSSARY OF LIMNOLOGICAL TERMS

**Aerobe-** Organisms requiring oxygen for life. All animals, most algae and some bacteria require oxygen for respiration.

**Algae-** See phytoplankton.

**Alkalinity-** Total concentration of bicarbonate and hydroxide ions (in most lakes).

**Anaerobe-** Organisms not requiring oxygen for life. Some algae and many bacteria are able to respire or ferment without using oxygen.

**Anoxic-** A system lacking oxygen, therefore incapable of supporting the most common kind of biological respiration, or of supporting oxygen-demanding chemical reactions. The deeper waters of a lake may become anoxic if there are many organisms depleting oxygen via respiration, and there is little or no replenishment of oxygen from photosynthesis or from the atmosphere.

**Benthic-** Referring to the bottom sediments.

**Bacterioplankton-** Bacteria adapted to the "open water" or "planktonic" zone of lakes, adapted for many specialized habitats and include groups that can use the sun's energy (phytoplankton), some that can use the energy locked in sulfur or iron, and others that gain energy by decomposing dead material.

**Bicarbonate-** The most important ion (chemical) involved in the buffering system of New Hampshire lakes.

**Buffering-** The capacity of lakewater to absorb acid with a minimal change in the pH. In New Hampshire the chemical responsible for buffering is the bicarbonate ion. (See pH.)

**Chloride-** One of the components of salts dissolved in lakewater. Generally the most abundant ion in New Hampshire lakewater, it may be used as an indicator of raw sewage or of road salt.

**Chlorophyll a-** The main green pigment in plants. The concentration of chlorophyll *a* in lakewater is often used as an indicator of algal abundance.

**Circulation-** The period during spring and fall when the combination of low water temperature and wind cause the water column to mix freely over its entire depth.

**Density-** The weight per volume of a substance. The more dense an object, the heavier it feels. Low-density liquids will float on higher-density liquids.

**Dimictic-** The thermal pattern of lakes where the lake circulates, or mixes, twice a year. Other patterns such as polymictic (many periods of circulation per year) are uncommon in New Hampshire. (See also meromictic and holomictic).

**Dystrophy-** The lake trophic state in which the lakewater is highly stained with humic acids (reddish brown or yellow stain) and has low productivity. Chlorophyll *a* concentration may be low or high.

**Epilimnion-** The uppermost layer of water during periods of thermal stratification. (See lake diagram).

**Eutrophy-** The lake trophic state in which algal production is high. Associated with eutrophy is low Secchi Disk depth, high chlorophyll *a*, and high total phosphorus. From an esthetic viewpoint these lakes are "bad" because water clarity is low, aquatic plants are often found in abundance, and cold-water fish such as trout and salmon are usually not present. A good aspect of eutrophic lakes is their high productivity in terms of warm-water fish such as bass, pickerel, and perch.

**Free CO<sub>2</sub>-** Carbon dioxide that is not combined chemically with lake water or any other substances. It is produced by respiration, and is used by plants and bacteria for photosynthesis.

**Holomixis-** The condition where the entire lake is free to circulate during periods of overturn. (See meromixis.)

**Humic Acids-** Dissolved organic compounds released from decomposition of plant leaves and stems. Humic acids are red, brown, or yellow in color and are present in nearly all lakes in New Hampshire. Humic acids are consumed only by fungi, and thus are relatively resistant to biological decomposition.

**Hydrogen Ion-** The "acid" ion, present in small amounts even in distilled water, but contributed to rain-water by atmospheric processes, to ground-water by soils, and to lakewater by biological organisms and sediments. The active component of "acid rain". See also "pH" the symbolic value inversely and exponentially related to the hydrogen ion.

**Hypolimnion-** The deepest layer of lakewater during periods of thermal stratification. (See lake diagram)

**Lake-** Any "inland" body of relatively "standing" water. Includes many synonyms such as ponds, tarns, loches, billabongs, bogs, marshes, etc.

**Lake Morphology-** The shape and size of a lake and its basin.

**Littoral-** The area of a lake shallow enough for submerged aquatic plants to grow.

**Meromixis-** The condition where the entire lake fails to circulate to its deepest points; caused by a high concentration of salt in the deeper waters, and by peculiar landscapes (small deep lakes surrounded by hills and/or forests. (Contrast holomixis.)

**Mesotrophy-** The lake trophic state intermediate between oligotrophy and eutrophy. Algal production is moderate, and chlorophyll  $\alpha$ , Secchi Disk depth, and total phosphorus are also moderate. These lakes are esthetically "fair" but not as good as oligotrophic lakes.

**Metalimnion-** The "middle" layer of the lake during periods of summer thermal stratification. Usually defined as the region where the water temperature changes at least one degree per meter depth. Also called the thermocline.

**Mixis-** Periods of lakewater mixing or circulation.

**Mixotrophy-** The lake condition where the water is highly stained with humic acids, but algal production and chlorophyll  $\alpha$  values are also high.

**Oligotrophy-** The lake trophic state where algal production is low, Secchi Disk depth is deep, and chlorophyll  $\alpha$  and total phosphorus are low. Esthetically these lakes are the "best" because they are clear and have a minimum of algae and aquatic plants. Deep oligotrophic lakes can usually support cold-water fish such as lake trout and land-locked salmon.

**Overturn-** See circulation or mixis

**pH-** A measure of the hydrogen ion concentration of a liquid. For every decrease of 1 pH unit, the hydrogen ion concentration increases 10 times. Symbolically, the pH value is the "negative logarithm" of the hydrogen ion concentration. For example, a pH of 5 represents a hydrogen ion concentration of  $10^{-5}$  molar. [Please thank the chemists for this lovely symbolism -- and ask them to explain it in lay terms!] In any event, the higher the pH value, the lower the hydrogen ion concentration. The range is 0 to 14, with 7 being neutral 1 denoting high acid condition and 14 denoting very basic condition.

**Photosynthesis-** The process by which plants convert the inorganic substances carbon dioxide and water into organic glucose (sugar) and oxygen using sunlight as the energy source. Glucose is an energy source for growth, reproduction, and maintenance of almost all life forms.

**Phytoplankton-** Microscopic algae which are suspended in the "open water" zone of lakes and ponds. A major source of food for zooplankton. Common examples include: diatoms, euglenoids, dinoflagellates, and many others. Usually included are the blue-green bacteria.

**Parts per million-** Also known as "ppm". This is a method of expressing the amount of one substance (solute) dissolved in another (solvent). For example, a solution with 10 ppm of oxygen has 10 pounds of oxygen for every 999,990 pounds (500 tons) of water. Domestic sewage usually contains from 2 to 10 ppm phosphorus.

**Parts per billion-** Also known as "ppb". This is only 1/1000 of ppm, therefore much less concentrated. As little as 1 ppb of phosphorus will sustain growth of algae. As little as 10 ppb phosphorus will cause algal blooms! Think of the ratio as 1 milligram (1/28000 of an ounce) of phosphorus in 25 barrels of water (55 gallon drums)! Or, 1 gallon of septic waste diluted into 10,000 gallons of lakewater. It adds up fast!



**Plankton**- Community of microorganisms that live suspended in the water column, not attached to the bottom sediments or aquatic plants. See also "bacterioplankton" (bacteria), "phytoplankton" (algae) and "zooplankton" (microcrustaceans and rotifers).

**Saturated**- When a solute (such as water) has dissolved all of a substance that it can. For example, if you add table salt to water, a point is reached where any additional salt fails to dissolve. The water is then said to be saturated with table salt. In lakewater, gaseous oxygen can dissolve, but eventually the water becomes saturated with oxygen if exposed sufficiently long to the atmosphere or another source of oxygen.

**Specific Conductivity**- A measure of the amount of salt present in lakewater. As the salt concentration increases, so does the specific conductivity (electrical conductivity).

**Stratum**- A layer or "blanket". Can be used to refer to one of the major layers of lakewater such as the epilimnion, or to any layers of organisms or chemicals that may be present in a lake.

**Thermal Stratification**- The process by which layers are built up in the lake due to heating by the sun and partial mixing by wind.

**Thermocline**- Region of temperature change. (See metalimnion.)

**Total Phosphorus**- A measure of the concentration of phosphorus in lakewater. Includes both free forms (dissolved), and chemically combined form (as in living tissue, or in dead but suspended organisms).

**Trophic Status**- A classification system placing lakes into similar groups according to their amount of algal production. (See Oligotrophy, Mesotrophy, Eutrophy, Mixotrophy, and Dystrophy for definitions of the major categories)

**Z**- A symbol used by limnologists as an abbreviation for depth.

**Zooplankton**- Microscopic animals in the planktonic community. Some are called "water fleas", but most are known by their scientific names. Scientific names include: *Daphnia*, *Cyclops*, *Bosmina*, and *Kellicottia*.